

# IRREDUCIBILITY AND SMOOTHNESS OF THE MODULI SPACE OF MATHEMATICAL 5-INSTANTONS OVER $\mathbb{P}_3$

I. COANDĂ<sup>1</sup>, A. TIKHOMIROV<sup>2</sup>, AND G. TRAUTMANN<sup>3</sup>

ABSTRACT. We prove that the space of mathematical instantons with second Chern class 5 over  $\mathbb{P}_3$  is smooth and irreducible. Unified and simple proofs for the same statements in case of second Chern class  $\leq 4$  are contained.

## CONTENTS

|                                              |    |
|----------------------------------------------|----|
| Introduction                                 | 1  |
| Notations                                    | 2  |
| 1. Instanton bundles                         | 3  |
| 2. Instanton Bundles and Multilinear Algebra | 5  |
| 3. The Method                                | 12 |
| 4. On Jumping Lines of $n$ -Instantons       | 18 |
| 5. The varieties $M(n, r)$ for $n \leq 4$    | 24 |
| 6. A technical result about 5-instantons     | 26 |
| 7. Irreducibility of $MI(5)$                 | 30 |
| 8. Smoothness of $MI(5)$                     | 42 |
| References                                   | 46 |

## INTRODUCTION

A mathematical instanton bundle is an algebraic vector bundle  $\mathcal{E}$  over  $\mathbb{P}_3(k)$ ,  $k$  an algebraically closed field of characteristic 0, if it has rank 2, Chern classes  $c_1 = 0$ ,  $c_2 = n > 0$ , and if it satisfies the vanishing conditions  $H^0\mathcal{E} = 0$  and  $H^1\mathcal{E}(-2) = 0$ . The name was chosen in twistor theory in the 1970's, when holomorphic bundles on  $\mathbb{P}_3(\mathbb{C})$  with the

---

<sup>1</sup> Research partially supported by the Romanian Academy Grant No. 60/2001 and by the EURROMAT programme sponsored by the European Commission.

<sup>2</sup>Research partially supported by the INTAS programme.

<sup>3</sup>Research partially supported by the DFG.

same properties were considered as counterparts of (anti-)self-dual Yang–Mills fields on the 4-sphere, see [1], [27], [12] for reference. Let a mathematical instanton bundle with  $c_2 = n$  be called  $n$ -instanton or  $(n, 2)$ -instanton for short. The isomorphism classes of  $n$ -instantons are the closed points of a coarse moduli scheme  $MI(n)$ . Since its first consideration, it is an open problem whether  $MI(n)$  is smooth and irreducible for any  $n$ . An affirmative answer for  $n \leq 4$  has been given in several papers, for each  $n$  separately, and recently Katsylo and Ottaviani proved smoothness for  $n = 5$ , see historical remarks in 1.3.

The main result of this paper is that also  $MI(5)$  is irreducible, with a new proof of smoothness included. The method used also enables a simple and unified proof for all the previous cases  $n \leq 4$ , see Section 5.

It is well-known that any  $n$ -instanton is the cohomology of a short complex  $H_n \otimes \mathcal{O}(-1) \rightarrow N \otimes \mathcal{O} \rightarrow H_n^* \otimes \mathcal{O}(1)$ , where  $H_n$  and  $N$  are  $k$ -vector spaces of dimensions  $n$  and  $2n + 2$ , respectively, also called the Horrocks construction. We consider also higher rank instanton bundles, called  $(n, r)$ -instantons, which can be constructed from the same type of complexes with the same  $H_n$  but with  $\dim N = 2n + r$ ,  $2 \leq r \leq 2n$ . We prove that, given an  $(n, r)$ -instanton  $\mathcal{E}$ , one can choose a linear form  $\xi \in H_n^*$  such that with  $H_{n-1} \cong \text{Ker}(\xi)$  the induced complex  $H_{n-1} \otimes \mathcal{O}(-1) \rightarrow N \otimes \mathcal{O} \rightarrow H_{n-1}^* \otimes \mathcal{O}(1)$  defines an  $(n-1, r+2)$ -instanton  $\mathcal{E}_\xi$ , together with a complex  $\mathcal{O}(-1) \rightarrow \mathcal{E}_\xi \rightarrow \mathcal{O}(1)$ , whose cohomology is the original  $\mathcal{E}$ , see Section 3. Together with technical details, this observation enables us to perform induction steps  $(n-1, r+2) \rightsquigarrow (n, r)$  for  $n \leq 5$  for irreducibility and smoothness. These induction steps are short for  $n \leq 4$ , see Section 5, while the case  $(4, 4) \rightsquigarrow (5, 2)$  is more elaborate. In Section 3, the induction step is explained in more details. The proof of smoothness in each case is achieved by proving that  $H^2 S^2 \mathcal{E}_\xi = 0$  implies  $H^2 S^2 \mathcal{E} = 0$  for suitable  $\xi$ .

## NOTATIONS

- Throughout the paper  $k$  will be an algebraically closed field of characteristic zero
- If  $E$  is a finite dimensional  $k$ -vector space,  $\mathbb{P}E$  will denote the projective space of 1-dimensional and  $G_m E$  the Grassmannian of  $m$ -dimensional subspaces.
- The invertible sheaf of degree  $d$  over  $\mathbb{P}E$  is denoted  $\mathcal{O}_{\mathbb{P}E}(d)$  such that  $E^* = H^0(\mathbb{P}E, \mathcal{O}_{\mathbb{P}E}(1))$ .
- $\mathbb{P} = \mathbb{P}V \cong \mathbb{P}_3$  will denote the projective 3-space over  $k$  for a fixed vector space  $V$  of dimension 4. We will omit the index  $\mathbb{P}$  at the structure sheaf  $\mathcal{O} = \mathcal{O}_{\mathbb{P}}$  and at the invertible sheaves  $\mathcal{O}(d) = \mathcal{O}_{\mathbb{P}}(d)$ .
- For an  $\mathcal{O}_{\mathbb{P}}$ -module  $\mathcal{F}$  we use the abbreviations  $\mathcal{F}(d) = \mathcal{F} \otimes \mathcal{O}_{\mathbb{P}}(d)$ ,  $H^i \mathcal{F} = H^i(\mathbb{P}, \mathcal{F})$  and  $h^i \mathcal{F}(d) = \dim_k H^i \mathcal{F}(d)$ , and  $\text{ext}^i(\mathcal{F}, \mathcal{G}) = \dim_k \text{Ext}^i(\mathcal{F}, \mathcal{G})$ .
- If  $E$  is a finite dimensional  $k$ -vector space, the sheaf of sections of the trivial vector bundle over a scheme  $X$  will be denoted  $E \otimes \mathcal{O}_X = E \otimes_k \mathcal{O}_X$  and  $E \otimes \mathcal{F}$  is written for  $(E \otimes \mathcal{O}_X) \otimes_{\mathcal{O}_X} \mathcal{F}$ .

- If  $\mathcal{F}$  respectively  $\mathcal{G}$  are coherent sheaves on schemes  $X$  resp.  $Y$ ,  $\mathcal{F} \boxtimes \mathcal{G}$  denotes the sheaf  $p^*\mathcal{F} \otimes q^*\mathcal{G}$  on  $X \times_k Y$  where  $p$  respectively  $q$  are the projections.
- If  $\mathcal{F}$  is a coherent sheaf on the scheme  $X$  and  $Y$  a closed subscheme of  $X$  we simply write  $\mathcal{F}_Y$  for the restriction on  $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_Y = \mathcal{F}|_Y$ .
- Throughout the paper a vector bundle is a locally free sheaf of finite rank. If necessary, we will refer specifically to its bundle space as a fibration of vector spaces.
- The Chern classes  $c_i(\mathcal{F})$  of a coherent sheaf  $\mathcal{F}$  over  $\mathbb{P} \cong \mathbb{P}_3$  are considered as integers and we also call the triple  $(c_1(\mathcal{F}), c_2(\mathcal{F}), c_3(\mathcal{F}))$  the Chern class of  $\mathcal{F}$ .

## 1. INSTANTON BUNDLES

**1.1.** Let  $M(2; 0, n, 0)$  denote the moduli space of semistable coherent sheaves on  $\mathbb{P}_3$  of rank 2 with Chern class  $(c_1, c_2, c_3) = (0, n, 0)$ , which had been constructed by M. Maruyama, [17], [18]. It contains the open set  $M^b(2; 0, n)$  of stable rank 2 vector bundles on  $\mathbb{P}_3$  with Chern class  $(c_1, c_2) = (0, n)$ . Recall that a rank 2 vector bundle  $\mathcal{E}$  on  $\mathbb{P}_3$  with Chern class  $c_1 = 0$  is stable if and only if  $H^0\mathcal{E} = 0$ . Then it is also simple, i.e.  $\text{hom}(\mathcal{E}, \mathcal{E}) = 1$ , see [20]. In this case the Riemann-Roch formula becomes

$$\text{ext}^1(\mathcal{E}, \mathcal{E}) - \text{ext}^2(\mathcal{E}, \mathcal{E}) = 8n - 3,$$

whereas  $\text{Ext}^1(\mathcal{E}, \mathcal{E})$  is isomorphic to the tangent space of  $M^b(2; 0, n)$  at  $[\mathcal{E}]$ , the isomorphism class of  $\mathcal{E}$ . For large  $n$  the spaces  $M^b(2; 0, n)$  have many irreducible components and some of them have a much bigger dimension than  $8n - 3$ .

**1.2. Definition:** A symplectic mathematical instanton bundle with second Chern class  $n \geq 0$  and of rank  $r$ , or an  $(n, r)$ -instanton for short, is a locally free sheaf  $\mathcal{E}$  over  $\mathbb{P} = \mathbb{P}V \cong \mathbb{P}_3$  with the following properties

- (i)  $\mathcal{E}$  has Chern class  $(0, n, 0)$  and  $2 \leq r = rk(\mathcal{E}) \leq 2n$
- (ii)  $H^0\mathcal{E} = 0$  and  $H^1\mathcal{E}(-2) = 0$
- (iii)  $\mathcal{E}$  admits a symplectic isomorphism  $\mathcal{E} \xrightarrow{\varphi} \mathcal{E}^*, \varphi^* = -\varphi$ .

Then  $\mathcal{E}$  must have even rank by (iii). In case  $rk(\mathcal{E}) = 2$  condition (iii) can be dropped because then the non-degenerate pairing  $\mathcal{E} \otimes \mathcal{E} \rightarrow \Lambda^2\mathcal{E} \cong \mathcal{O}$  determines a symplectic form, which then is unique up to a scalar.

If  $rk(\mathcal{E}) = 2$ , then  $\mathcal{E}$  is stable by condition (ii). Therefore, the open part

$$MI(n) \subset M^b(2; 0, n)$$

defined by the condition  $h^1\mathcal{E}(-2) = 0$  is the set of isomorphism classes of  $(n, 2)$ -instantons, also called  $n$ -instantons.

**1.3. Conjecture:**  $MI(n)$  is smooth and irreducible of dimension  $8n - 3$  for  $n \geq 1$ .

This conjecture emerged in the late 1970's as  $n$ -instantons were considered as counterparts of self-dual Yang–Mills fields on the 4-sphere, see [1], [27], in twistor theory.

The conjecture has been proved for  $n \leq 4$ . For  $n = 1$  the space  $MI(1)$  is the complement of the Grassmannian  $G(2, V)$  in  $\mathbb{P}(\Lambda^2 V)$ . For  $n = 2$  the space  $MI(2)$  had been described by R. Hartshorne in [12] as a smooth and irreducible fibration. The case  $n = 3$  was proved by G. Ellingsrud–S.A. Strømme in [10]. W. Barth proved in [3] that  $MI(4)$  is irreducible and J. LePotier in [21] that  $MI(4)$  is smooth. Recently P.I. Katsylo and G. Ottaviani proved that also  $MI(5)$  is smooth, see [15]. In this paper we prove that  $MI(5)$  is both smooth and irreducible. The method also yields simple and unified proofs for the previous results for  $n \leq 4$ .

#### 1.4. Further results on $n$ -instantons:

(1) Any  $n$ -instanton  $\mathcal{E}$  is stable because  $H^0\mathcal{E} = 0$  and  $\text{rk}(\mathcal{E}) = 2$ . Then the Grauert–Mülich theorem states that  $\mathcal{E}$  has trivial splitting type, i.e. for a general line  $L$  in  $\mathbb{P}_3$  the restricted bundle  $\mathcal{E}_L$  is isomorphic to  $2\mathcal{O}_L$ , see [2].

(2) A line  $L$  is called a jumping line of the  $n$ -instanton  $\mathcal{E}$  if  $\mathcal{E}_L \cong \mathcal{O}_L(-a) \oplus \mathcal{O}_L(a)$  with  $a \neq 0$ , and this number is called the order of the jumping line. It is an easy consequence of the monad representation of  $\mathcal{E}$ , see 2.5 that the set  $J(\mathcal{E})$  of all jumping lines of  $\mathcal{E}$  is a hypersurface of degree  $n$  in the Grassmannian  $\mathbb{G}$  of lines in  $P_3$ . Moreover,  $n$  is the highest order possible for a jumping line.

(3) For any  $n$ -instanton,  $h^0\mathcal{E}(1) \leq 2$ , see [6]. The  $n$ -instantons with  $h^0\mathcal{E}(1) = 2$  are called special 'tHooft bundles. They can be presented as extensions

$$0 \rightarrow 2\mathcal{O}_{\mathbb{P}_3}(-1) \rightarrow \mathcal{E} \rightarrow \mathcal{O}_Q(-n, 1) \rightarrow 0,$$

where  $Q$  is a smooth quadric in  $\mathbb{P}_3$ , see [13]. It was shown there that  $MI(n)$  is smooth along the locus of special 'tHooft bundles, which has dimension  $2n + 9$ .

(4) A plane  $P$  in  $\mathbb{P}_3$  is called unstable for an  $n$ -instanton  $\mathcal{E}$  if the restricted bundle  $\mathcal{E}_P$  has sections, otherwise stable. If  $\mathcal{E}$  is special 'tHooft, the unstable planes form a smooth quadric surface in  $\mathbb{P}_3^*$ , the dual of  $Q$  in (3). In [7] it was proved that an  $n$ -instanton  $\mathcal{E}$  is already special 'tHooft if its variety of unstable planes has dimension  $\geq 2$ .

(5) In [19] it was shown that  $MI(n)$  is smooth at any  $\mathcal{E}$  with  $h^0\mathcal{E}(1) \neq 0$  and that  $\text{Ext}^2(\mathcal{E}, \mathcal{E}(-1)) = 0$  for such a bundle. The locus of these bundles has dimension  $5n + 4$ .

(6) P. Rao, [22], and M. Skiti, [24], proved independently that  $MI(n)$  is even smooth along the locus of bundles  $\mathcal{E}$  which admit jumping lines of highest order  $n$ . These bundles form a subvariety of dimension  $6n + 2$ . Moreover, in [22] it is proved that  $MI(5)$  is smooth at bundles which have jumping lines of order 4.

(7) L. Costa and G. Ottaviani, [9], proved that  $MI(5)$  is an affine scheme by describing the non-degeneracy condition for the monads, see 2.2, as the non-vanishing of a hyperdeterminant.

## 2. INSTANTON BUNDLES AND MULTILINEAR ALGEBRA

In this section we describe the monad construction of instantons from hypernets of quadrics. Throughout this paper

$$H = H_n$$

denotes an  $n$ -dimensional  $k$ -vector space,  $n \geq 1$ . We identify  $\Lambda^2(H^* \otimes V^*)$  with the space of anti-selfdual  $k$ -linear maps

$$H \otimes V \longrightarrow H^* \otimes V^*.$$

Each  $\omega \in \Lambda^2(H^* \otimes V^*)$  gives rise to a diagram

$$\begin{array}{ccccccc} H \otimes V & \xrightarrow{\omega} & H^* \otimes V^* & \longrightarrow & Q & \longrightarrow & 0 \\ \downarrow u^* & & \uparrow u & & & & \\ N^* & \xrightarrow[\sim]{\varphi} & N & & & & \end{array}$$

where  $N = N_\omega$  is the image and  $Q = Q_\omega$  the cokernel of  $\omega$ , and  $\varphi = \varphi_\omega$  is the canonically induced symplectic isomorphism. Note that the rank of  $\omega$  is always even.  $\omega$  is called **non-degenerate** if  $\omega(h \otimes v) \neq 0$  for any non-zero decomposable tensor  $h \otimes v$  in  $H \otimes V$ .

**2.1. Rank Stratification:** The space  $\Lambda^2(H^* \otimes V^*)$  comes with the rank stratification

$$\Lambda^2(H^* \otimes V^*) = \Omega_{4n} \supset \Omega_{4n-2} \supset \dots \supset \Omega_2,$$

where  $\Omega_{2m} = \Omega_{2m}(H) = \{\omega \in \Lambda^2(H^* \otimes V^*) \mid \text{rk}(\omega) \leq 2m\}$ . This had been considered already in [25] by A. Tyurin.

It is easy to prove by standard arguments that each  $\Omega_{2m}$  is irreducible and smooth outside  $\Omega_{2m-2}$  of codimension  $\binom{4n-2m}{2}$ . The tangent space at  $\omega \in \Omega_{2m} \setminus \Omega_{2m-2}$  is the kernel in

$$0 \rightarrow T_\omega \Omega_{2m} \rightarrow \Lambda^2(H^* \otimes V^*) \rightarrow \Lambda^2 Q \rightarrow 0$$

which gives the dimension formula. There is the canonical decomposition

$$\Lambda^2(H^* \otimes V^*) = (S^2 H^* \otimes \Lambda^2 V^*) \oplus (\Lambda^2 H^* \otimes S^2 V^*),$$

and on the first summand we have the (induced) rank stratification

$$S^2 H^* \otimes \Lambda^2 V^* = M_{4n} \supset M_{4n-2} \supset \dots \supset M_2$$

with  $M_{2m} = M_{2m}(H) = \Omega_{2m} \cap (S^2 H^* \otimes \Lambda^2 V^*)$ . We let

$$\Delta = \Delta(H) = \{\omega \in S^2 H^* \otimes \Lambda^2 V^* \mid \omega \text{ is degenerate}\}$$

be the subset of degenerate tensors.

**Remark:** One can prove that  $\Delta$  is a closed and irreducible subvariety of  $S^2 H^* \otimes \Lambda^2 V^*$  of codimension  $2(n-1)$  for  $n \geq 2$  and of codimension 1 for  $n = 1$ .

**2.2. The bundles  $\mathcal{E}_\omega$ .** To each  $\omega \in S^2 H^* \otimes \Lambda^2 V^*$  one can associate the complex

$$H \otimes \mathcal{O}_{\mathbb{P}}(-1) \xrightarrow{\varphi \circ \alpha^*} N \otimes \mathcal{O}_{\mathbb{P}} \xrightarrow{\alpha} H^* \otimes \mathcal{O}_{\mathbb{P}}(1),$$

where  $\alpha$  is the composition of  $N \otimes \mathcal{O}_{\mathbb{P}} \xrightarrow{u} H^* \otimes V^* \otimes \mathcal{O}_{\mathbb{P}}$  and  $H^* \otimes V^* \otimes \mathcal{O}_{\mathbb{P}} \xrightarrow{\text{id} \otimes \text{ev}} H^* \otimes \mathcal{O}_{\mathbb{P}}(1)$ . Then the following are equivalent:

- (i)  $\alpha$  is surjective
- (ii)  $\varphi \circ \alpha^*$  is a subbundle
- (iii)  $\omega$  is non-degenerate.

Clearly (i) and (ii) are equivalent by definition. Now  $\alpha$  is surjective if and only if for any  $\langle v \rangle \in \mathbb{P}V$  the induced homomorphism  $N \rightarrow H^* \otimes \langle v \rangle^*$  on the fibre is surjective or equivalently, that  $H \otimes \langle v \rangle \xrightarrow{\omega} H^* \otimes V^*$  is injective, which is the non-degeneracy.

By this observation each  $\omega \in M_{2m} \setminus (M_{2m-2} \cup \Delta)$ ,  $n < m \leq 2n$ , gives rise to an associated locally free sheaf

$$\mathcal{E}_\omega = \text{Ker}(\alpha) / \text{Im}(\varphi \circ \alpha^*).$$

This bundle is in fact an instanton bundle of rank  $2m - 2n$  and with Chern class  $(c_1, c_2, c_3) = (0, n, 0)$ . This follows directly from the defining complex, also called the **monad** of  $\mathcal{E}_\omega$ . Rank and Chern classes can be computed from those of the sheaves of the complex as well as  $H^0 \mathcal{E}_\omega = 0$  and  $H^1 \mathcal{E}_\omega(-2) = 0$ . Because  $\mathcal{E}_\omega^*$  is the cohomology of the dual complex, it follows that the symplectic isomorphism  $N^* \xrightarrow{\varphi} N$  induces a symplectic isomorphism  $\mathcal{E}_\omega^* \xrightarrow{j_\omega} \mathcal{E}_\omega$ .

**Remark:**  $M_{2n} \subset \Delta$  because otherwise a non-degenerate  $\omega \in M_{2n} \setminus M_{2n-2}$  would define the bundle  $\mathcal{E}_\omega = 0$  with non-trivial second Chern class. Therefore,  $2n + 2$  is the lowest possible rank for a tensor  $\omega \in S^2 H^* \otimes \Lambda^2 V^*$ , which is non-degenerate, and which then defines an instanton of rank 2.

In the following lemma and proposition it is proved that any  $(n, r)$ -instanton arises by the above construction and that the isomorphism classes of  $(n, r)$ -instantons are in 1 : 1 correspondence with the equivalence classes of the operators  $\omega \in M_{2n+r} \setminus (M_{2n+r-2} \cup \Delta)$ . We use the following **notation** for  $2 \leq r \leq 2n$ .

$$M(n, r) := M_{2n+r} \setminus (M_{2n+r-2} \cup \Delta)$$

or

$$M(n, r) := \{\omega \in S^2 H_n^* \otimes \Lambda^2 V^* \mid rk(\omega) = 2n + r, \ \omega \text{ non-degenerate}\}.$$

**2.2.1. Lemma:** *For any  $\omega \in M(n, r)$ , there are isomorphisms*

$$\begin{aligned} H &\cong H^2(\mathcal{E}_\omega \otimes \Omega^3(1)) & N &\cong H^1(\mathcal{E}_\omega \otimes \Omega^1) & Q &\cong H^1 \mathcal{E}_\omega \\ H^* &\cong H^1 \mathcal{E}_\omega(-1) & N^* &\cong H^2(\mathcal{E}_\omega \otimes \Omega^2) \end{aligned}$$

*which are compatible with Serre duality and the symplectic isomorphism  $\mathcal{E}_\omega^* \cong \mathcal{E}_\omega$ , making the following diagram commute*

$$\begin{array}{ccccccccc} H \otimes V & \longrightarrow & N^* & \xrightarrow[\sim]{\varphi} & N & \longrightarrow & H^* \otimes V^* & \longrightarrow & Q \longrightarrow 0 \\ \downarrow \approx & & \downarrow \approx & & \downarrow \approx & & \downarrow \approx & & \downarrow \approx \\ H^2(\mathcal{E}_\omega \otimes \Omega^3(1)) \otimes V & \longrightarrow & H^2(\mathcal{E}_\omega \otimes \Omega^2) & \xleftarrow[\sim]{\delta} & H^1(\mathcal{E}_\omega \otimes \Omega^1) & \longrightarrow & H^1(\mathcal{E}_\omega(-1)) \otimes V^* & \longrightarrow & H^1 \mathcal{E}_\omega \longrightarrow 0 \end{array}$$

*Proof.* The homomorphisms of the bottom row of the diagram are induced by the natural homomorphism  $\Omega^3(1) \otimes V \rightarrow \Omega^2$  and the exact sequences  $0 \rightarrow \Omega^2 \rightarrow \Lambda^2 V^* \otimes \mathcal{O}(-2) \rightarrow \Omega^1 \rightarrow 0$  and  $0 \rightarrow \Omega^1 \rightarrow V^* \otimes \mathcal{O}(-1) \rightarrow \mathcal{O} \rightarrow 0$  of the Koszul complex of  $V^* \otimes \mathcal{O}(-1) \rightarrow \mathcal{O}$ . Then the isomorphisms of the lemma and the commutativity of the diagram follow by tracing all data from the defining complex of  $\mathcal{E}_\omega$ , using the functoriality of Serre-duality. See also [4] for monads of vector bundles on projective spaces.  $\square$

**2.3. Proposition:** (a) *Let  $\mathcal{E}$  be an  $(n, r)$ -instanton with symplectic isomorphism  $\mathcal{E}^* \xrightarrow{j} \mathcal{E}$ ,  $2 \leq r \leq 2n$ . Then there exists a non-degenerate  $\omega \in S^2 H^* \otimes \Lambda^2 V^*$  of rank  $2n + r$  such that  $(\mathcal{E}, j) \cong (\mathcal{E}_\omega, j_\omega)$ .*

(b)  *$(\mathcal{E}_\omega, j_\omega) \cong (\mathcal{E}_{\omega'}, j_{\omega'})$  if and only if  $\omega$  and  $\omega'$  are in the same  $GL(H)$ -orbit in  $S^2 H^* \otimes \Lambda^2 V^*$ .*

(c) *The stabilizer  $Stab(\omega)$  of  $\omega$  in  $GL(H)$  is isomorphic to the automorphism group  $Aut(\mathcal{E}_\omega, j_\omega)$ .*

*Proof.* (a) By 2.2.1 it is enough to show that the Beilinson spectral sequence of  $\mathcal{E}$  results in the complex

$$0 \rightarrow H^2(\mathcal{E} \otimes \Omega^3(1)) \otimes \mathcal{O}(-1) \xrightarrow{\beta} H^1(\mathcal{E} \otimes \Omega^1) \otimes \mathcal{O} \xrightarrow{\alpha} H^1(\mathcal{E}(-1)) \otimes \mathcal{O}(1) \rightarrow 0,$$

where  $\alpha$  resp.  $\beta$  correspond to the natural homomorphisms

$$H^2(\mathcal{E} \otimes \Omega^3(1)) \otimes V \rightarrow H^2(\mathcal{E} \otimes \Omega^2) \xleftrightarrow[\sim]{\sim} H^1(\mathcal{E} \otimes \Omega^1)$$

resp.

$$H^1(\mathcal{E} \otimes \Omega^1) \rightarrow H^1 \mathcal{E}(-1) \otimes V^*,$$

which are Serre–dual to each other. Now for any coherent sheaf  $\mathcal{F}$  on  $\mathbb{P}_n$  there is the Beilinson I complex

$$0 \rightarrow C^{-n}(\mathcal{F}) \rightarrow \cdots \rightarrow C^0(\mathcal{F}) \rightarrow \cdots \rightarrow C^n(\mathcal{F}) \rightarrow 0$$

with terms  $C^p(\mathcal{F}) = \bigoplus_{p=i-j} H^i(\mathcal{F} \otimes \Omega^j(j)) \otimes \mathcal{O}(-j)$ , which is exact except at  $C^0(\mathcal{F})$  and which has  $\mathcal{F}$  as its cohomology at  $C^0(\mathcal{F})$ . In our case for  $\mathcal{F} = \mathcal{E}(-1)$  this complex reduces in fact to

$$0 \rightarrow H^1(\mathcal{E} \otimes \Omega^2(1)) \otimes \mathcal{O}(-2) \rightarrow H^1(\mathcal{E} \otimes \Omega^1) \otimes \mathcal{O}(-1) \rightarrow H^1(\mathcal{E}(-1)) \otimes \mathcal{O} \rightarrow 0$$

by verifying that the instanton conditions imply the vanishing of the other terms of the Beilinson complex. Moreover, the Koszul sequence  $0 \rightarrow \Omega^3 \rightarrow \Lambda^3 V^* \otimes \mathcal{O}(-3) \rightarrow \Omega^2 \rightarrow 0$  induces the isomorphism  $H^1(\mathcal{E} \otimes \Omega^2(1)) \cong H^2(\mathcal{E} \otimes \Omega^3(1))$  because  $H^i \mathcal{E}(-2) = 0$  for  $i = 1, 2$ . Now  $\mathcal{E}$  is the cohomology of the complex

$$0 \rightarrow H^2(\mathcal{E} \otimes \Omega^3(1)) \otimes \mathcal{O}(-1) \xrightarrow{\beta} H^1(\mathcal{E} \otimes \Omega^1) \otimes \mathcal{O} \xrightarrow{\alpha} H^1(\mathcal{E}(-1)) \otimes \mathcal{O}(1) \rightarrow 0.$$

It follows from the Riemann–Roch formula that

$$\chi \mathcal{E}(d) = r \binom{d+3}{3} - n(d+2)$$

and from the instanton conditions that

$$h^1 \mathcal{E}(-1) = n \quad \text{and} \quad h^1(\mathcal{E} \otimes \Omega^1) = 2n + r.$$

Thus  $\mathcal{E}$  defines a non-degenerate  $\omega \in S^2 H^* \otimes \Lambda^2 V^*$  via

$$H^2(\mathcal{E} \otimes \Omega^3(1)) \otimes V \rightarrow H^2(\mathcal{E} \otimes \Omega^2) \xleftarrow{\sim} H^1(\mathcal{E} \otimes \Omega^1) \rightarrow H^1 \mathcal{E}(-1) \otimes V^*$$

together with an isomorphism  $H^* \cong H^1 \mathcal{E}(-1)$ , such that  $(\mathcal{E}, j) \cong (\mathcal{E}_\omega, j_\omega)$ .

(b) and (c) follow from (a) and 2.2.1 by the isomorphisms of the complexes.  $\square$

**2.4. Corollary:** *If  $\mathcal{E}_\omega$  is simple, then  $\text{Stab}_{GL(H)}(\omega) = \{\pm id_H\}$ .*

**2.5. Remark:** There is also a Beilinson II monad for an instanton bundle  $\mathcal{E}_\omega$ . This can be treated in the same way as the above Beilinson I monad. Any  $\omega \in S^2 H^* \otimes \Lambda^2 V^*$  defines an operator  $H \rightarrow H^* \otimes \Lambda^2 V^*$  which is symmetric with respect to  $H$  and the exact sequence  $0 \rightarrow N \xrightarrow{u} H^* \otimes V^* \rightarrow Q \rightarrow 0$  as above. Now combined with the Koszul homomorphisms

$$\Lambda^2 V^* \otimes \mathcal{O}(-1) \rightarrow \Omega^1(1) \quad \text{and} \quad \Omega^1(1) \rightarrow V^* \otimes \mathcal{O}$$

we obtain the complex

$$0 \rightarrow H \otimes \mathcal{O}(-1) \xrightarrow{\mu} H^* \otimes \Omega^1(1) \xrightarrow{\nu} Q \otimes \mathcal{O} \rightarrow 0.$$



If  $\omega$  is non-degenerate, this complex defines the  $(n, r)$ -instanton  $\mathcal{E}_\omega$ , where  $\text{rk}(\omega) = 2n + r$ . This could be shown as in 2.2.1, 2.3, but follows already from the following commutative diagram induced by  $\omega$ , which is a direct transformation between the two monad types.

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & H \otimes \mathcal{O}(-1) & \xrightarrow{\mu} & H^* \otimes \Omega^1(1) & \longrightarrow & Q \otimes \mathcal{O} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & N \otimes \mathcal{O} & \xrightarrow{u} & H^* \otimes V^* \otimes \mathcal{O} & \longrightarrow & Q \otimes \mathcal{O} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & H^* \otimes \mathcal{O}(1) & \xlongequal{\quad} & H^* \otimes \mathcal{O}(1) & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 
 \end{array}$$

## 2.6. The morphism $M_{2n+2} \setminus \Delta \xrightarrow{b} MI(n)$

According to Proposition 2.3 we are given a surjective map  $\omega \xrightarrow{b} [\mathcal{E}_\omega]$  from  $M_{2n+2} \setminus \Delta$  to  $MI(n)$  whose fibres are the orbits under the action of  $GL(H)$ . The map  $b$  is the underlying map of a morphism, because there is a universal monad over  $(M_{2n+2} \setminus \Delta) \times \mathbb{P}$  with the universal family of  $n$ -instantons as cohomology. Then  $b$  is the modular morphism onto the open part  $MI(n)$  of the Maruyama scheme. It can be shown that  $b$  is a geometric quotient and a principal  $GL(H)/\{\pm 1\}$ -bundle in the etale topology. However the latter fact will not be used in this paper.

## 2.7. Criterion for irreducibility

Since  $\Omega_{2n+2}$  is irreducible of codimension  $\binom{2n-2}{2}$  in  $\Lambda^2(H^* \otimes V^*)$ , it follows that every irreducible component of  $M_{2n+2}$  has codimension  $\leq \binom{2n-2}{2}$  in  $S^2 H^* \otimes \Lambda^2 V^*$  or dimension  $\geq 3n(n+1) - \binom{2n-2}{2} = n^2 + 8n - 3$ . From this observation we obtain

**2.7.1. Lemma:** *Any component of  $MI(n)$  has dimension  $\geq 8n - 3$  and  $MI(n)$  is irreducible of dimension  $8n - 3$  if and only if  $M_{2n+2} \setminus \Delta$  is irreducible of the expected codimension  $\binom{2n-2}{2}$  in  $S^2 H^* \otimes \Lambda^2 V^*$ .*

For the proof use the following lemma.

**2.7.2. Lemma:** *Let  $X \xrightarrow{f} Y$  be a morphism of reduced schemes, let  $Y$  be irreducible and assume that each fibre  $f^{-1}(f(x))$ ,  $x \in X$ , is irreducible of constant dimension  $e$ , and that each irreducible component of  $X$  is of dimension  $\geq \dim Y + e$ . Then  $X$  is irreducible.*

*Proof.* Let  $X_1, \dots, X_p$  be the irreducible components of  $X$ . According to the theorem of the dimension of the fibres,  $f|_{X_i}$  is dominant over  $Y$  for any  $i$ . Let  $X'_i \subset X_i$  be the complement of the other components and let  $y \in \cap f(X'_i)$ . Then

$$f^{-1}(y) = \cup (f^{-1}(y) \cap X_i)$$

and the intersections  $f^{-1}(y) \cap X_i$  are distinct closed subsets of  $f^{-1}(y)$ . As  $f^{-1}(y)$  is irreducible, it follows that  $p = 1$ .  $\square$

**2.7.3. Remark:** In [26], Example 1, A. Tyurin showed that  $M_{2n}$  and (consequently)  $M_{2n+2}$  have not the expected codimension  $\binom{2n}{2}$  respectively  $\binom{2n-2}{2}$  in  $S^2 H^* \otimes \Lambda^2 V^*$ .

## 2.8. Criterion for smoothness

Concerning the smoothness of  $MI(n)$ , let  $\mathcal{E}$  be an  $n$ -instanton (of rank 2). Because  $[\mathcal{E}]$  is a stable point of  $MI(n)$ , this is a smooth point if  $\text{Ext}^2(\mathcal{E}, \mathcal{E}) = 0$ . But since  $\text{ext}^1(\mathcal{E}, \mathcal{E}) - \text{ext}^2(\mathcal{E}, \mathcal{E}) = 8n - 3$  the vanishing of  $\text{Ext}^2(\mathcal{E}, \mathcal{E})$  is also necessary if  $[\mathcal{E}]$  is a smooth point and  $MI(n)$  has dimension  $8n - 3$  at  $[\mathcal{E}]$ . On the other hand, we have

$$\text{Ext}^2(\mathcal{E}, \mathcal{E}) \cong H^2(\mathcal{E}^\vee \otimes \mathcal{E}) \cong H^2(\mathcal{E} \otimes \mathcal{E}) \cong H^2(S^2 \mathcal{E}).$$

Now for any  $(n, r)$ -instanton we have the following

**2.8.1. Lemma:** *Let  $\omega \in S^2 H^* \otimes \Lambda^2 V^*$  be non-degenerate of rank  $2m$ ,  $n < m \leq 2n$ , and let  $\mathcal{E} = \mathcal{E}_\omega$ . Then the following conditions are equivalent.*

- (i)  $H^2(S^2 \mathcal{E}) = 0$
- (ii)  $\Omega_{2m}$  and  $S^2 H^* \otimes \Lambda^2 V^*$  intersect transversally in  $\omega$  inside  $\Lambda^2(H^* \otimes V^*)$ .
- (iii)  $M_{2m}$  is smooth at  $\omega$  of expected dimension  $5n - 5n^2 + 8mn - 2m^2 - m$ .

*Proof.* Transversality in (ii) means that the tangent space  $T_\omega \Omega_{2m}$  and  $S^2 H^* \otimes \Lambda^2 V^*$  span  $\Lambda^2(H^* \otimes V^*)$ . Then (ii) and (iii) are equivalent by standard dimension counts, because  $\Omega_{2m}$  has codimension  $\binom{4n-2m}{2}$  and is smooth at  $\omega$ .

Next, let us recall a general fact. To any bounded complex  $K^\bullet$  of vector bundles on a variety one can associate its second symmetric power  $S^2 K^\bullet$  by decomposing  $K^\bullet \otimes K^\bullet$  according to the eigenspaces of the canonical involution on  $K^\bullet \otimes K^\bullet$ . If  $K^\bullet$  has only three non-zero terms  $K^{-1} \rightarrow K^0 \rightarrow K^1$ , then  $S^2 K^\bullet$  is isomorphic to

$$0 \rightarrow \Lambda^2 K^{-1} \rightarrow K^{-1} \otimes K^0 \rightarrow S^2 K^0 \oplus (K^{-1} \otimes K^1) \rightarrow K^0 \otimes K^1 \rightarrow \Lambda^2 K^1 \rightarrow 0$$

whose differentials are naturally deduced from those of  $K^\bullet$ . Moreover, if  $H^i(K^\bullet) = 0$  for  $i \neq 0$  and  $H^0(K^\bullet) = F$ , then  $H^i(S^2K^\bullet) = 0$  for  $i \neq 0$  and  $H^0(S^2K^\bullet) \cong S^2F$ . Hence, in our case  $S^2\mathcal{E}$  is the degree 0 cohomology of the derived complex

$$\Lambda^2 H \otimes \mathcal{O}(-2) \rightarrow H \otimes N \otimes \mathcal{O}(-1) \rightarrow (S^2 N \oplus H \otimes H^*) \otimes \mathcal{O} \rightarrow N \otimes H^* \otimes \mathcal{O}(1) \rightarrow \Lambda^2 H^* \otimes \mathcal{O}(2).$$

The terms of this monad are cohomologically acyclic. Hence one can compute the cohomology of  $S^2\mathcal{E}$  just by passing to global sections and then taking cohomology. Particularly, one gets an exact sequence

$$N \otimes H^* \otimes V^* \rightarrow \Lambda^2 H^* \otimes S^2 V^* \rightarrow H^2(S^2\mathcal{E}) \rightarrow 0.$$

Now condition (ii) is equivalent to

$$T_\omega \Omega_{2m} + (S^2 H^* \otimes \Lambda^2 V^*) = \Lambda^2(H^* \otimes V^*),$$

noting that  $\omega$  is a smooth point of  $\Omega_{2m}$ . Taking into account the exact sequences

$$N \otimes (H^* \otimes V^*) \rightarrow \Lambda^2(H^* \otimes V^*) \rightarrow \Lambda^2 Q \rightarrow 0$$

and

$$0 \rightarrow S^2 H^* \otimes \Lambda^2 V^* \rightarrow \Lambda^2(H^* \otimes V^*) \rightarrow \Lambda^2 H^* \otimes S^2 V^* \rightarrow 0,$$

one finds that (ii) is equivalent to the surjectivity of the composed map

$$N \otimes H^* \otimes V^* \rightarrow \Lambda^2(H^* \otimes V^*) \rightarrow \Lambda^2 H^* \otimes S^2 V^*,$$

which is the map whose cokernel is  $H^2(S^2\mathcal{E})$ . This proves the equivalence of (i) and (ii).  $\square$

Now, by the criteria for irreducibility and smoothness, it is clear that the conjecture 1.3 on  $MI(n)$  is equivalent to

**2.9. Transcribed Conjecture:**  $\Omega_{2n+2}$  and  $S^2 H^* \otimes \Lambda^2 V^*$  intersect transversally inside  $\Lambda^2(H^* \otimes V^*)$  along  $M_{2n+2} \setminus \Delta$  and this intersection is irreducible.

Note, that in this conjecture, with  $\Delta$  also  $M_{2n}$  has been extracted from  $M_{2n+2}$  because  $M_{2n} \subset \Delta$ . According to Tyurin's example the whole of  $M_{2n+2}$  has components of excessive dimension for large  $n$ .

**2.10. Remark:** Katsylo–Ottaviani gave in [15] the following interpretation of the transversality condition. Dualizing the sequence with  $H^2(S^2\mathcal{E})$  as cokernel in the proof of Lemma 2.8.1, we have the exact sequence

$$0 \rightarrow H^2(S^2\mathcal{E})^* \rightarrow \Lambda^2 H \otimes S^2 V \rightarrow N^* \otimes H \otimes V.$$

Interpreting the elements of  $\Lambda^2 H \otimes S^2 V$  as anti-selfdual linear maps  $H^* \otimes V^* \rightarrow H \otimes V$  and the elements of  $N^* \otimes H \otimes V$  as linear maps  $N \rightarrow H \otimes V$ , the morphism in the above sequence can be described by  $\sigma \mapsto \sigma \circ u$ , where  $u$  is the inclusion of  $\text{Im}(\omega) = N \subset H^* \otimes V^*$ . Therefore,

$$H^2(S^2 \mathcal{E})^* \cong \{\sigma \in \Lambda^2 H \otimes S^2 V \mid \sigma \circ \omega = 0\}$$

and the three conditions of 2.8.1 are equivalent to

$$(iv) \text{ if } \sigma \circ \omega = 0 \quad \text{for } \sigma \in \Lambda^2 H \otimes S^2 V, \quad \text{then } \sigma = 0.$$

In order to illustrate this point of view and for later use, we prove

**2.11. Proposition:**  $M(n, 2n - 2) = M_{4n-2} \setminus (\Delta \cup M_{4n-4})$  is smooth and (obviously) of codimension 1 in  $S^2 H^* \otimes \Lambda^2 V^*$  for  $n \geq 2$ .

*Proof.* Let  $\omega \in M_{4n-2} \setminus (\Delta \cup M_{4n-4})$  and  $\sigma \in \Lambda^2 H \otimes S^2 V$  such that  $\sigma \circ \omega = 0$ . Since both  $\omega$  and  $\sigma$  are anti-selfdual, it follows that  $\omega \circ \sigma = 0$ . Now  $\text{rk}(\omega) = 4n - 2$  implies  $\text{rk}(\sigma) \leq 2$ . One can now easily show (see e.g. [25], Proposition 2.1.1) that in this case  $\sigma$  is decomposable,  $\sigma = \eta \otimes f$ , with  $\eta \in \Lambda^2 H$  of rank  $\leq 2$  and  $f \in S^2 V$  of rank  $\leq 1$ . Consequently, if  $\sigma \neq 0$ , then  $\text{Im}(\sigma) \subset H \otimes V$  contains decomposable vectors, contradicting  $\omega \notin \Delta$ .  $\square$

### 3. THE METHOD

Let  $\mathcal{E}$  be an  $(n, r)$ -instanton and  $H^* = H_n^* \cong H^1 \mathcal{E}_\omega(-1)$  as above with  $\omega \in S^2 H^* \otimes \Lambda^2 V^* \setminus \Delta$  of rank  $2n + r$ . To any  $\xi \in H^*$  with kernel  $\bar{H}$  we can associate the restriction  $\bar{\omega} = \omega_\xi = \text{res}_\xi(\omega)$  of  $\omega$  to  $S^2 \bar{H}^* \otimes \Lambda^2 V^*$  such that we have the diagram

$$\begin{array}{ccc} \bar{H} \otimes V & \xrightarrow{\bar{\omega}} & \bar{H}^* \otimes V^* \\ \downarrow & & \uparrow \\ H \otimes V & \xrightarrow{\omega} & H^* \otimes V^* \end{array}$$

Then  $\text{rk}(\bar{\omega}) \leq \text{rk}(\omega)$ .

The choice of  $\xi$  gives rise to a diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \bar{H} \otimes \mathcal{O}(-1) & & \mathcal{O}(1) & & \\
 & & \downarrow & \searrow \bar{\beta} & \downarrow & & \\
 0 & \longrightarrow & H \otimes \mathcal{O}(-1) & \xrightarrow{\beta} & N \otimes \mathcal{O} & \xrightarrow{\alpha} & H^* \otimes \mathcal{O}(1) \longrightarrow 0 \\
 & & \downarrow & & \searrow \bar{\alpha} & & \downarrow \\
 & & \mathcal{O}(-1) & & \bar{H}^* \otimes \mathcal{O}(1) & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 
 \end{array}$$

such that  $\bar{\beta}, \bar{\alpha}$  constitute a monad for a bundle  $\bar{\mathcal{E}}$  with an induced symplectic isomorphism, and such that  $\bar{\beta}, \bar{\alpha}$  are induced by  $\bar{\omega}$ . Then  $\bar{\mathcal{E}}$  has rank  $r + 2$  and second Chern class  $n - 1$ . Any splitting homomorphism of  $H \otimes \mathcal{O}(-1) \rightarrow \mathcal{O}(-1)$  then induces a unique homomorphism  $\mathcal{O}(-1) \rightarrow \bar{\mathcal{E}}$  and by duality a monad

$$0 \rightarrow \mathcal{O}(-1) \rightarrow \bar{\mathcal{E}} \rightarrow \mathcal{O}(1) \rightarrow 0$$

whose cohomology is again  $\mathcal{E}$ . This observation will be used to perform an induction  $(n - 1, r + 2) \rightsquigarrow (n, r)$  for irreducibility and smoothness of the spaces of  $(n, r)$ -instantons for  $n \leq 5$ .

Immediate relations between the tensors  $\omega$  and  $\bar{\omega}$  and the corresponding sheaves  $\mathcal{E}$  and  $\bar{\mathcal{E}}$  are stated in the next two lemmata.

**3.1. Lemma:** *Let  $\bar{\omega}$  be the restriction of  $\omega$  for  $\xi \in H^*$ . The following conditions are equivalent:*

- (i)  $rk(\bar{\omega}) = rk(\omega)$
- (ii)  $Im(\omega) \cap (\xi \otimes V^*) = 0$
- (iii) *The multiplication map  $H^1\mathcal{E}(-1) \otimes V^* \rightarrow H^1\mathcal{E}$  restricts to an injection  $\xi \otimes V^* \hookrightarrow H^1\mathcal{E}$ .*
- (iv)  $H^0\bar{\mathcal{E}} = 0$

*Proof.* We have the diagram

$$\begin{array}{ccccccc}
 \bar{H} \otimes V & \xrightarrow{\bar{\omega}} & \bar{H}^* \otimes V^* & & & & \\
 \downarrow j & & \uparrow j^* & & & & \\
 H \otimes V & \xrightarrow{\omega} & H^* \otimes V^* & \xrightarrow{\mu} & Q & \longrightarrow & 0 \\
 \downarrow u^* & & \uparrow u & & & & \\
 N^* & \xrightarrow[\approx]{\varphi} & N & & & & 
 \end{array}$$

where  $N$  is the image of  $\omega$ ,  $\text{Ker}(j^*) = \xi \otimes V^*$ , and  $\mu$  is isomorphic to the multiplication map  $H^1\mathcal{E}(-1) \otimes V^* \rightarrow H^1\mathcal{E}$ . It follows that  $j^* \circ u$  is injective if and only if the image of  $\bar{\omega}$  has the same dimension as  $N$ , which proves the equivalence of (i) and (ii). The other statements are immediately seen to be equivalent to the injectivity of  $j^* \circ u$ . For (iv), note, that  $N \rightarrow \bar{H}^* \otimes V^*$  corresponds to the right part of the monad of  $\bar{\mathcal{E}}$ .

□

If the conditions of the lemma are satisfied, then  $\bar{\mathcal{E}}$  is an  $(n-1, r+2)$ -instanton, provided  $r < 2n$ . In this case we have

**3.2. Lemma:** *Let  $\omega \in S^2H^* \otimes \Lambda^2V^*$  be non-degenerate and  $\text{rk}(\bar{\omega}) = \text{rk}(\omega)$ . If  $H^2(S^2\bar{\mathcal{E}}) = 0$  and  $H^1\mathcal{E}(1) = 0$ , then  $H^2(S^2\mathcal{E}) = 0$ .*

*Proof.* Because  $\mathcal{E}$  is the cohomology of the monad  $\mathcal{O}(-1) \rightarrow \bar{\mathcal{E}} \rightarrow \mathcal{O}(1)$ ,  $S^2\mathcal{E}$  is the cohomology of the induced monad, see proof of Lemma 2.8.1,

$$\bar{\mathcal{E}}(-1) \rightarrow \mathcal{O} \oplus S^2\bar{\mathcal{E}} \rightarrow \bar{\mathcal{E}}(1).$$

The computation of  $H^2(S^2\mathcal{E})$  from this monad gives the vanishing.

□

### 3.3. Construction lemma

For  $n \geq 3$  and  $2 \leq r \leq 2n-2$  and for any  $\xi \in H^*$  we consider the open subset

$$M(n, r)_\xi := \{\omega \in M(n, r) \mid \text{Im}(\omega) \cap (\xi \otimes V^*) = 0\}.$$

This is exactly the open subset of  $M(n, r)$  which is mapped to  $M(n-1, r+2)$  under the restriction map

$$\text{res}_\xi : \omega \mapsto \omega_\xi = \omega \mid S^2\bar{H}^* \otimes \Lambda^2V^*.$$

We also use the notation

$$M^0(n, r) = \{\omega \in M(n, r) \mid H^1\mathcal{E}_\omega(1) = 0\}.$$

Note that this set might be empty if  $h^0\mathcal{E}_\omega(1) - h^1\mathcal{E}_\omega(1) = 4r - 3n < 0$ , while  $h^2\mathcal{E}_\omega(1) = h^3\mathcal{E}_\omega(1) = 0$ . Next we let

$$M(n, r)_\xi' = \{\omega \in M(n, r)_\xi \mid \text{res}_\xi(\omega) \in M^0(n-1, r+2)\}$$

be the inverse image. For the fibres of  $\text{res}_\xi$  we have the following lemma, which enables the induction. Note, that in this lemma the fibre  $R(\bar{\omega})$  may contain degenerate  $\omega$ 's.

**3.3.1. Lemma:** *a) For any  $\bar{\omega} \in M(n-1, r+2)$  there is an isomorphism*

$$R(\bar{\omega}) := \{\omega \in S^2 H_n^* \otimes \Lambda^2 V^* \mid \text{res}_\xi(\omega) = \bar{\omega}, \quad \text{rk}(\omega) = \text{rk}(\bar{\omega})\} \cong \text{Hom}_{\mathcal{O}_{\mathbb{P}}}(\mathcal{C}_{\bar{\omega}}, \mathcal{O}_{\mathbb{P}}(1))$$

where  $\mathcal{C}_{\bar{\omega}}$  denotes the cokernel of the left part of the monad of  $\mathcal{E}_{\bar{\omega}}$ , and

$$\dim \text{Hom}_{\mathcal{O}_{\mathbb{P}}}(\mathcal{C}_{\bar{\omega}}, \mathcal{O}_{\mathbb{P}}(1)) = n - 1 + h^0\mathcal{E}_{\bar{\omega}}(1).$$

*b)  $\omega \in R(\bar{\omega})$  is non-degenerate if and only if the corresponding homomorphism induces an epimorphism  $\mathcal{E}_{\bar{\omega}} \rightarrow \mathcal{O}_{\mathbb{P}}(1)$ . In that case  $\omega$  defines an  $(n, r)$ -instanton bundle  $\mathcal{E}_\omega$  which is also the cohomology of the self-dual monad  $\mathcal{O}_{\mathbb{P}}(-1) \rightarrow \mathcal{E}_{\bar{\omega}} \rightarrow \mathcal{O}_{\mathbb{P}}(1)$  defined by the epimorphism.*

*Proof.* a) For  $\xi$  fixed we may choose a decomposition  $H = H_1 \oplus \bar{H}$  such that  $\xi$  induces an isomorphism  $H_1 \cong k$ . Let  $\bar{\omega}$  decompose into

$$\bar{H} \otimes V \xrightarrow{\bar{u}^*} N^* \xrightarrow[\approx]{\varphi} N \xrightarrow{\bar{u}} \bar{H}^* \otimes V^*$$

with associated monad

$$\bar{H} \otimes \mathcal{O}_{\mathbb{P}}(-1) \xrightarrow{\bar{\beta}} N \otimes \mathcal{O}_{\mathbb{P}} \xrightarrow{\bar{\alpha}} \bar{H}^* \otimes \mathcal{O}_{\mathbb{P}}(1).$$

With  $\mathcal{C}_{\bar{\omega}} = \text{coker}(\bar{\beta})$  we have the exact sequence

$$0 \rightarrow \mathcal{E}_{\bar{\omega}} \rightarrow \mathcal{C}_{\bar{\omega}} \rightarrow \bar{H}^* \otimes \mathcal{O}_{\mathbb{P}}(1) \rightarrow 0.$$

Any  $u_1 \in \text{Hom}(N, H_1^* \otimes V^*)$  gives rise to a skew-symmetric operator

$$\omega = \left( \begin{array}{c|c} u_1 \circ \varphi \circ u_1^* & u_1 \circ \varphi \circ \bar{u}^* \\ \hline \bar{u} \circ \varphi \circ u_1^* & \bar{\omega} \end{array} \right)$$

$$(H_1 \oplus \bar{H}) \otimes V \xrightarrow{\omega} (H_1^* \oplus \bar{H}^*) \otimes V^*$$

which factors also through  $\varphi$  by its definition, so that  $\text{rk}(\omega) = \text{rk}(\bar{\omega})$  and  $\omega$  is symplectic. However, the component  $u_1 \circ \varphi \circ \bar{u}^* : \bar{H} \otimes V \rightarrow H_1^* \otimes V^*$  is not necessarily skew with respect to  $V$ . This is the case if and only if  $\omega \in R(\bar{\omega})$  or if and only if the composition

$$\bar{H} \otimes \mathcal{O}_{\mathbb{P}}(-1) \xrightarrow{\bar{\beta}} N \otimes \mathcal{O}_{\mathbb{P}} \xrightarrow{\bar{u}_1} H_1^* \otimes \mathcal{O}_{\mathbb{P}}(1)$$

of the associated sheaf homomorphism is zero, or, if and only if  $\tilde{u}_1$  factors through  $\mathcal{C}_{\bar{\omega}}$ . Let  $\text{Hom}(N, H_1^* \otimes V^*)'$  denote the subspace of  $\text{Hom}(N, H_1^* \otimes V^*)$  defined by this condition. Then  $u_1 \mapsto \omega$  defines an isomorphism

$$\text{Hom}(N, H_1^* \otimes V^*)' \cong R(\bar{\omega})$$

because  $u_1 \mapsto \omega$  is injective since  $\varphi$  and  $\bar{u}$  are surjective and because any  $\omega \in R(\bar{\omega})$  arises in this way. On the other hand, the factorization of  $\tilde{u}_1 : N \otimes \mathcal{O}_{\mathbb{P}} \twoheadrightarrow \mathcal{C}_{\bar{\omega}} \xrightarrow{\gamma} H_1^* \otimes \mathcal{O}_{\mathbb{P}}(1)$  defines an isomorphism  $u_1 \leftrightarrow \gamma$  between  $\text{Hom}(N, H_1^* \otimes V^*)'$  and  $\text{Hom}(\mathcal{C}_{\bar{\omega}}, H_1^* \otimes \mathcal{O}_{\mathbb{P}}(1))$ .

b) By a) any  $\omega \in R(\bar{\omega})$  gives rise to a diagram

$$\begin{array}{ccccccc}
 & & & & 0 & & \\
 & & & & \uparrow & & \\
 & & & & \bar{H}^* \otimes \mathcal{O}_{\mathbb{P}}(1) & & \\
 & & \nearrow \bar{\alpha} & & \uparrow & & \\
 0 \longrightarrow & \bar{H} \otimes \mathcal{O}_{\mathbb{P}}(-1) & \xrightarrow{\bar{\beta}} & N \otimes \mathcal{O}_{\mathbb{P}} & \longrightarrow & \mathcal{C}_{\bar{\omega}} & \longrightarrow 0 \\
 & \downarrow \alpha_1 & & \nwarrow & & \uparrow & \\
 & H_1^* \otimes \mathcal{O}_{\mathbb{P}}(1) & & \longleftarrow \pi & & \mathcal{E}_{\bar{\omega}} & \\
 & & & & & \uparrow & \\
 & & & & & 0 & 
 \end{array}$$

with  $\pi$  induced by  $u_1$  or  $\alpha_1$ . It follows that  $\alpha = \alpha_1 + \bar{\alpha} : N \otimes \mathcal{O}_{\mathbb{P}} \rightarrow (H_1^* \oplus \bar{H}^*) \otimes \mathcal{O}_{\mathbb{P}}(1)$  is surjective if and only if  $\pi$  is surjective. This proves b), because the surjectivity of  $\alpha$  (as right part of the monad of  $\omega$ ) is equivalent to the non-degeneracy of  $\omega$ . In that case the induced sequence

$$H_1 \otimes \mathcal{O}_{\mathbb{P}}(-1) \rightarrow \mathcal{E}_{\bar{\omega}} \xrightarrow{\pi} H_1^* \otimes \mathcal{O}_{\mathbb{P}}(1)$$

is also a monad for the bundle  $\mathcal{E}_{\omega}$ . □

**3.3.2. Corollary:** *If  $M^0(n-1, r+2)$  is irreducible of the expected dimension  $3(n-1)n - \binom{2n-4-r}{2}$ , then also  $M(n, r)'_{\xi}$  is irreducible of the expected dimension*

$$3n(n+1) - \binom{2n-r}{2},$$

*if it is not empty.*

*Proof.* The dimension of any component of  $M(n, r)$  is  $\geq$  then the expected dimension by 2.7. Because the fibres of  $\text{res}_{\xi}$  have constant dimension by 3.3.1, the corollary follows from Lemma 2.7.2. □



**3.4. Lemma:** *If  $M(n, r)'_\xi \neq \emptyset$ , then for any other  $\eta \neq 0$ ,*

$$M(n, r)'_\xi \cap M(n, r)'_\eta \neq \emptyset.$$

*Proof.* For  $\omega_0 \in M(n, r)'_\xi$  we consider the set

$$U_0 = \{\eta \in H^* \mid \text{Im}(\omega_0) \cap (\eta \otimes V^*) = 0, h^1 \mathcal{E}_\eta(1) = 0\}$$

where  $\mathcal{E}_\eta$  denotes the bundle obtained from  $\omega^0$  by restriction to the kernel of  $\eta$ . Because  $(\mathcal{E}_\eta)$  is a flat family on the open set of  $\eta$ 's defined by  $\text{Im}(\omega_0) \cap (\eta \otimes V^*) = 0$ , the semicontinuity theorem implies that also  $U_0$  is open. Because  $\xi \in U_0$ , there is a  $\xi' \in U_0$  which is independent of  $\xi$ . Then  $\omega_0 \in M(n, r)'_\xi \cap M(n, r)'_{\xi'}$ .

Let now  $\eta \neq 0$  be arbitrary in  $H^*$ . There is a transformation  $g \in GL(H)$  such that  $g^* \xi = \xi$  and  $g^* \xi' = \eta$ . Then

$$(g^* \otimes \text{id}_{V^*}) \circ \omega_0 \circ (g \otimes \text{id}_V) \in M(n, r)'_\xi \cap M(n, r)'_\eta.$$

□

**3.5. The induction:** Suppose now that  $M^0(n-1, r+2)$  is irreducible and that also  $M(n, r)'_\xi \neq \emptyset$  for some  $\xi \neq 0$ . Then by Lemma 3.4 the union

$$\bigcup_{\xi \neq 0} M(n, r)'_\xi$$

is an irreducible open subset of  $M(n, r)$  of the expected dimension. If, in addition,  $M^0(n-1, r+2)$  is smooth (as transversal intersection), then by Lemma 3.2 also  $H^2 S^2 \mathcal{E}_\omega = 0$  for any  $\omega$  in the above union. Concerning rank 2 instantons, we shall prove that

$$M(n, 2) = \bigcup_{\xi \neq 0} M(n, 2)_\xi$$

for  $n \geq 3$ . For  $n = 3, 4$  we shall even prove that

$$M(n, 2) = \bigcup_{\xi \neq 0} M(n, 2)'_\xi.$$

This will give a unified proof of the instanton conjecture for  $n \leq 4$ .

For  $n = 5$ , however, we are at present not able to prove that for any  $\omega \in M(5, 2)$  there exists a  $\xi \neq 0$  such that  $H^1 \mathcal{E}_\omega(1) = 0$ . But we shall prove the weaker result, that there exists a  $\xi \neq 0$  with  $h^1 \mathcal{E}_\omega(1) \leq 1$ . This is already sufficient to prove the conjecture for  $n = 5$  in the sequel.

For  $n = 6$  one might hope that  $\bigcup_{\xi} M(6, 2)'_\xi$  is the complement of the subvariety of 'tHooft bundles (defined by  $h^0 \mathcal{E}(1) \neq 0$ ). However, for  $n > 6$  the present induction method doesn't seem to work anymore.

4. ON JUMPING LINES OF  $n$ -INSTANTONS

In this section we are going to prove the following Proposition 4.5 which enables us to choose  $\xi \in H_n^*$  for any  $n$ -instanton  $\mathcal{E}_\omega, n \geq 3$ , such that  $\mathcal{E}_\xi := \mathcal{E}_\omega$  is again an  $(n-1, 4)$ -instanton, i.e.  $H^0 \mathcal{E}_\xi = 0$ . For  $n \geq 5$  the proposition allows us to choose a second  $\eta \in H_{n-1}^*$  such that also  $H^0(\mathcal{E}_\xi)_\eta = 0$

In order to prepare the proof we include the following lemmata on jumping lines of arbitrary instantons.

**4.1. Lemma:** *Let  $\mathcal{E}$  be a stable rank 2 vector bundle on  $\mathbb{P}_2$  with Chern class  $(c_1, c_2) = (0, n)$ ,  $n \geq 2$ , let  $L_1, L_2$  be distinct lines and let  $a_1, a_2 \geq 0$  be defined by  $\mathcal{E}_{L_\nu} \cong \mathcal{O}_{L_\nu}(a_\nu) \oplus \mathcal{O}_{L_\nu}(-a_\nu)$ . Then*

(a)  $a_1 + a_2 \leq n$

(b) *If  $a_1 + a_2 = n$ , and  $a_1 \geq 2, a_2 \geq 2$ , then  $\mathcal{E}$  can be realized as an extension*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_2}(-1) \rightarrow \mathcal{E} \rightarrow \mathcal{I}_Z(1) \rightarrow 0$$

*with  $Z \subset L_1 \cup L_2$ , a 0-dimensional subscheme of length  $n+1$ .*

*Proof.* (a) There is the natural exact sequence

$$0 \rightarrow \mathcal{O}_{L_1 \cup L_2} \rightarrow \mathcal{O}_{L_1} \oplus \mathcal{O}_{L_2} \rightarrow \mathcal{O}_{L_1 \cap L_2} \rightarrow 0. \quad (1)$$

Tensoring this with  $\mathcal{E}$  and taking sections, this implies

$$a_1 + a_2 \leq h^0 \mathcal{E}_{L_1 \cup L_2}.$$

On the other hand, the sequence

$$0 \rightarrow \mathcal{E}(-2) \rightarrow \mathcal{E} \rightarrow \mathcal{E}_{L_1 \cup L_2} \rightarrow 0 \quad (2)$$

implies the exact sequence

$$0 \rightarrow H^0 \mathcal{E} \rightarrow H^0 \mathcal{E}_{L_1 \cup L_2} \rightarrow H^1 \mathcal{E}(-2) \rightarrow H^1 \mathcal{E}.$$

Because  $\mathcal{E}$  is stable,  $h^0 \mathcal{E} = 0$  and  $h^2 \mathcal{E}(-2) = h^0 \mathcal{E}(-1) = 0$ , so that  $h^1 \mathcal{E}(-2) = n$  by the Riemann–Roch formula, and then  $h^0 \mathcal{E}_{L_1 \cup L_2} \leq n$ .

(b) According to the proof of (a), if  $a_1 + a_2 = n$ , then  $a_1 + a_2 = h^0 \mathcal{E}_{L_1 \cup L_2} = n$  and

$$H^0 \mathcal{E}_{L_1} \oplus H^0 \mathcal{E}_{L_2} \rightarrow H^0 \mathcal{E}_{L_1 \cap L_2}$$

is surjective. It follows that then also

$$H^0 \mathcal{E}_{L_1}(1) \oplus H^0 \mathcal{E}_{L_2}(1) \rightarrow H^0 \mathcal{E}_{L_1 \cap L_2}(1)$$

is surjective and from this, that

$$h^0 \mathcal{E}_{L_1 \cup L_2}(1) = n + 2.$$

Using sequence (2) with  $\mathcal{E}(1)$ , we obtain  $h^0 \mathcal{E}(1) \geq 2$ . Then there are exact sequences

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_2}(-1) \rightarrow \mathcal{E} \rightarrow \mathcal{I}_Z(1) \rightarrow 0$$

with 0-dimensional subschemes  $Z$  of  $\mathbb{P}_2$  of length  $n+1$ . Tensoring (1) with  $\mathcal{O}_Z$  and taking lengths, one obtains

$$\begin{aligned} \text{length}(Z \cap (L_1 \cup L_2)) &\geq \text{length}(Z \cap L_1) + \text{length}(Z \cap L_2) - 1 \\ &= (a_1 + 1) + (a_2 + 1) - 1 \\ &= n + 1 \\ &= \text{length}(Z). \end{aligned}$$

Then  $Z \subset L_1 \cup L_2$  as schemes.  $\square$

**4.2. Lemma:** *Let  $\mathcal{E}$  be a semistable rank 2 vector bundle on  $\mathbb{P}_2$  with Chern class  $(c_1, c_2) = (0, n)$ ,  $n \geq 2$ . Then*

- (a) *For odd  $n \geq 3$  or even  $n \geq 6$ ,  $\mathcal{E}$  has only finitely many jumping lines of order  $\geq n/2$ .*
- (b) *For  $n = 2$  or 4,  $\mathcal{E}$  has at most one jumping line of order  $> n/2$ .*

*Proof.* Note, that semistability in this case means that  $H^0\mathcal{E}(-1) = 0$ , see [20]. Let firstly  $\mathcal{E}$  be stable with  $H^0\mathcal{E} = 0$ . Then 4.1, (a), implies that  $\mathcal{E}$  has at most one jumping line of order  $> n/2$ . Suppose  $n \geq 4$  is even. According to 4.1, (b), if  $\mathcal{E}$  has two jumping lines  $L_1, L_2$  of order  $\geq n/2$ , then there is an extension

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_2}(-1) \rightarrow \mathcal{E} \rightarrow \mathcal{I}_Z(1) \rightarrow 0$$

with  $Z \subset L_1 \cup L_2$  of length  $n+1$ . If  $L$  is any other line, then

$$\text{length}(Z \cap L) \leq \text{length}(L \cap (L_1 \cup L_2)) = 2,$$

and it follows that  $\mathcal{E}_L \cong \mathcal{O}_L(a) \oplus \mathcal{O}_L(-a)$  with  $0 \leq a \leq 1$ .

Now assume that  $\mathcal{E}$  is properly semistable. In this case we have an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_2} \rightarrow \mathcal{E} \rightarrow \mathcal{I}_Z \rightarrow 0$$

with a 0-dimensional scheme  $Z$  of length  $n$ . If  $L$  is a jumping line of  $\mathcal{E}$  order  $\geq n/2$ , then also  $\text{length}(Z \cap L) \geq n/2$  by the exact sequence tensored with  $\mathcal{O}_L$ . If  $\mathcal{E}$  should have infinitely many jumping lines of order  $\geq n/2$ , then there is a point  $x \in Z$  such that infinitely many such jumping lines meet  $Z$  exactly in  $x$ . Then  $\text{length}(\mathcal{O}_{Z,x} \otimes \mathcal{O}_{L,x}) \geq n/2$  for infinitely many lines. It follows that the germ  $(Z, x)$  is defined by two equations  $f, g \in \mathcal{O}_{\mathbb{P}_2, x}$  ( $Z$  is a locally complete intersection) with  $\text{mult}(f), \text{mult}(g) \geq n/2$ . Then

$$n = \text{length}(Z) \geq \text{length}(\mathcal{O}_{Z,x}) \geq \text{mult}(f)\text{mult}(g).$$

This is not possible for odd  $n \geq 3$  or even  $n \geq 6$ . For  $n = 2$  or 4 the same kind of argument as in the last part of the proof of 4.1, (b), shows that  $\mathcal{E}$  has at most one jumping line of order  $> n/2$ .  $\square$

**4.3. Proposition:** *Let  $\mathcal{E}$  be an  $n$ -instanton on  $\mathbb{P}_3$ ,  $n \geq 2$ . Then the set of jumping lines of  $\mathcal{E}$  of order  $> n/2$  has dimension  $\leq 1$ .*

*Proof.* Let  $\mathbb{G}$  denote the Grassmannian of lines in  $\mathbb{P}_3$  and consider the incidence diagram

$$\mathbb{G} \xleftarrow{p} \mathbb{F} \xrightarrow{q} \mathbb{P}_3^*$$

of lines in planes. Let  $\Sigma$  be the set of jumping lines of order  $> n/2$ . According to 4.2 the projection  $q|_{p^{-1}(\Sigma)}$  has finite fibres. Suppose  $\dim \Sigma \geq 2$ . Then  $q(p^{-1}(\Sigma)) = \mathbb{P}_3^*$ . Let then  $P_0$  be a stable plane for  $\mathcal{E}$ . It must contain a jumping line  $L_0$  of order  $> n/2$ . For a general plane  $P$  containing  $L_0$ ,  $\mathcal{E}_P$  is stable and, according to 4.1, (a),  $L_0$  is the only jumping line of order  $> n/2$  contained in  $P$ . On the other hand, for any  $P$  containing  $L_0$  the bundle  $\mathcal{E}_P$  is semistable. Hence, by 4.2,  $P$  contains at most finitely many jumping lines of order  $> n/2$ . Consequently, there are only finitely many jumping lines  $L$  of order  $> n/2$  meeting  $L_0$ . This means that  $T_{L_0}(\mathbb{G}) \cap \Sigma$  is a finite set, where  $T_{L_0}(\mathbb{G})$  denotes the geometric tangent hyperplane to  $\mathbb{G}$  at  $L_0$  in  $\mathbb{P}_5$ , whose intersection with  $\mathbb{G}$  is the cone of lines meeting  $L_0$ . This contradicts  $\dim \Sigma \geq 2$ .  $\square$

**4.4. Remark:** One can show, using the method of R. Strano and M. Green, as in [8], that a non special 'tHooft  $n$ -instanton  $\mathcal{E}$ ,  $n \geq 5$ , satisfies  $h^0 \mathcal{E}_P(1) \leq 1$  for a general plane  $P$ . Then, using the above arguments, one deduces that the set of jumping lines of  $\mathcal{E}$  of order  $\geq n/2$  is at most 1-dimensional. But this improves 4.2 only for even  $n \geq 6$ .

From the above statements on jumping lines we can now derive the following proposition, which is the key of the induction process of this paper.

**4.5. Proposition:** *a) Let  $\mathcal{E}$  be any  $n$ -instanton. If  $n \geq 3$ , then for a general  $\xi \in H^1 \mathcal{E}(-1)$  the multiplication map  $\xi \otimes V^* \rightarrow H^1 \mathcal{E}$  is injective.*

*b) If  $n \geq 5$ , then for a general 2-dimensional subspace  $U \subset H^1 \mathcal{E}(-1)$  the multiplication map  $U \otimes V^* \rightarrow H^1 \mathcal{E}$  is injective.*

Note, that the condition in a) means that  $\text{Im}(\omega) \cap (\xi \otimes V^*) = 0$ , see 3.1, because the multiplication map is isomorphic to  $H \otimes V^* \rightarrow Q$ .

*Proof.* a) The right part  $N \otimes \mathcal{O} \rightarrow H^* \otimes \mathcal{O}(1)$  of the monad of  $\mathcal{E}$  corresponds to the exact sequence  $0 \rightarrow N \rightarrow H^* \otimes V^* \xrightarrow{\mu} Q \rightarrow 0$ , where  $\mu$  is isomorphic to the multiplication map. If  $P \subset \mathbb{P}$  is a plane with equation  $z \in V^*$ , we obtain the exact sequence

$$0 \rightarrow H^0 \mathcal{E}_P \rightarrow N \rightarrow H^* \otimes V^* / \langle z \rangle$$

and, therefore,

$$H^0 \mathcal{E}_P \cong N \cap (H^* \otimes z). \tag{1}$$

On the other hand, if  $h^0 \mathcal{E}_P \neq 0$ , we are given an exact sequence

$$0 \rightarrow \mathcal{O}_P \rightarrow \mathcal{E}_P \rightarrow \mathcal{I}_Z \rightarrow 0$$

where  $Z$  is a 0-dimensional subscheme of  $P$  of length  $n$ , because  $h^0 \mathcal{E}_P(-1) = 0$ . It follows that

$$0 \leq h^0 \mathcal{E}_P \leq 1. \quad (2)$$

We let now  $\mathbb{P}(N)_1$  denote the set of decomposable classes  $\langle \xi \otimes z \rangle \in \mathbb{P}(H^* \otimes V^*)$  which are contained in  $P(N)$ , so that

$$\mathbb{P}(N)_1 = \mathbb{P}(N) \cap S_1$$

where  $S_1$  is the image of the Segre embedding  $\mathbb{P}H^* \times \mathbb{P}V^* \subset \mathbb{P}(H^* \otimes V^*)$ . There are the two projections

$$\mathbb{P}H^* \xleftarrow{p_1} \mathbb{P}(N)_1 \xrightarrow{q_1} \mathbb{P}V^*.$$

The isomorphism (1) implies that for any  $z \neq 0$  we have

$$\mathbb{P}(H^0 \mathcal{E}_P) \cong q_1^{-1}(\langle z \rangle),$$

where  $P$  is the plane with equation  $z$ . Therefore, the image of  $q_1$  is contained in the subvariety  $JP(\mathcal{E}) \subset \mathbb{P}V^*$  of unstable planes of  $\mathcal{E}$ .

Now (2) implies that  $\mathbb{P}(N)_1$  is isomorphic to  $JP(\mathcal{E})$ . Because  $\dim JP(\mathcal{E}) \leq 2$ , see [2], it follows that  $p_1(\mathbb{P}(N)_1)$  has dimensions  $\leq 2$ . This proves a) in case  $n \geq 4$ . If  $n = 3$  and  $\mathcal{E}$  is not special 'tHooft, then  $\dim JP(\mathcal{E}) \leq 1$  by 1.4, (4), so that a) is true in that case, too. But in case  $n = 3$  and  $\mathcal{E}$  is special 'tHooft, the claim follows from remark 4.6 below.

b) For the proof of part b) we consider the intersection

$$\mathbb{P}(N)_2 = \mathbb{P}(N) \cap S_2$$

with the secant variety of  $S_1$  in  $\mathbb{P}(H^* \otimes V^*)$ , such that  $\mathbb{P}(N)_2$  consists of all elements of type  $\langle \xi_1 \otimes z_1 + \xi_2 \otimes z_2 \rangle$  which are contained in  $\mathbb{P}(N)$ . Then we have the two projections

$$G(2, H^*) \xleftarrow{p_2} \mathbb{P}(N)_2 \setminus \mathbb{P}(N)_1 \xrightarrow{q_2} G(2, V^*).$$

We have to show that a general 2-dimensional subspace  $K \subset H^*$  satisfies  $N \cap (K \otimes V^*) = 0$ .

Now  $N \cap (K \otimes V^*) \neq 0$  if and only if either  $\mathbb{P}(K) \cap p_1(\mathbb{P}(N)_1) \neq \emptyset$  or  $K \in \text{Im}(p_2)$ .

Because  $\dim p_1(\mathbb{P}(N)) \leq 2$  by part a), it follows that the set of  $K$  with  $\mathbb{P}(K) \cap p_1(\mathbb{P}(N)_1) \neq \emptyset$  has dimension  $\leq 2 + (n - 2) = n$ . By assumption,  $\dim G(2, H^*) = 2(n - 2) > n + 1$  for  $n \geq 6$  and  $= n + 1$  for  $n = 5$ . Therefore, it remains to prove that  $\dim \text{Im}(p_2) \leq n + 1$  for  $n \geq 6$  and  $\dim \text{Im}(p_2) \leq 5$  for  $n = 5$ .

In order to derive the estimates, we consider the fibres of  $q_2$ . Let  $W \subset V^*$  be any 2-dimensional subspace with the dual line  $L \subset \mathbb{P}V$ . Then

$$N \cap (H^* \otimes W) \cong H^0 \mathcal{E}_L$$

as can easily be derived from the monad description of  $\mathcal{E}$ . On the other hand,

$$q_2^{-1}(W) = \mathbb{P}(N \cap H^* \otimes W) \setminus \mathbb{P}(N)_1$$

by definition of  $\mathbb{P}(N)_2$ . Therefore,

$$\dim q_2^{-1}(W) = h^0 \mathcal{E}_L - 1. \quad (3)$$

Let  $G(2, V^*) \supset J_1 \supset J_2 \supset \cdots \supset J_n$  be the filtration by the sets  $J_k$  of jumping lines  $L$  of order  $h^0 \mathcal{E}_L - 1 \geq k$ . It follows that

$$\dim q_2^{-1}(J_k \setminus J_{k+1}) \leq 3 + k \leq n$$

for  $k \leq n - 3$ . For  $k = n - 2$

$$\dim q_2^{-1}(J_{n-2} \setminus J_{n-1}) \leq 1 + (n - 2) = n - 1$$

because of Lemma 4.3 since  $n - 2 > n/2$ , and finally

$$\dim q_2^{-1}(J_{n-1} \setminus J_n) \leq n, \quad \dim q_2^{-1}(J_n) \leq n + 1.$$

Totally we have  $\dim(\mathbb{P}(N)_2 \setminus \mathbb{P}(N)_1) \leq n + 1$ , which is sufficient for  $n \geq 6$ . In case  $n = 5$ ,  $\dim(\mathbb{P}(N)_2 \setminus \mathbb{P}(N)_1) = 6$  is only possible if  $\dim J_5 = 1$ . In that case  $\mathcal{E}$  is a special 'tHooft bundle, and it follows by the direct argument in remark 4.6, that there are 2-dimensional subspaces  $K \subset H^*$  with  $N \cap (K \otimes V^*) = 0$ . This proves b).  $\square$

**4.6. Remark:** Let  $\mathcal{E}$  be an  $(n, 2)$ -instanton on  $\mathbb{P}_3$  with  $n \geq 2m + 2 \geq 3$ . One can prove, along the same lines, that then, for a general  $m$ -dimensional subspace  $K \subset H^1 \mathcal{E}(-1)$  the multiplication map  $K \otimes V^* \rightarrow H^1 \mathcal{E}$  is injective. If  $\mathcal{E}$  is a special 'tHooft bundle, this can be verified by the following direct argument. Let  $z_0, \dots, z_3$  be a basis of  $V^*$ . Then bases of  $N$  and  $H^*$  can be chosen such that the operator  $N \rightarrow H^* \otimes V^*$ , i.e. the right part of the monad of  $\mathcal{E}$ , can be represented by the matrix

$$\begin{pmatrix} z_0 & z_1 & z_2 & z_3 & & & \\ & & z_0 & z_1 & z_2 & z_3 & \\ & & & \ddots & & & \ddots \\ & & & & z_0 & z_1 & z_2 & z_3 \end{pmatrix}.$$

If  $n = 2m + 1$  or  $2m + 2$  and  $e_1, \dots, e_n$  is the corresponding basis of  $H^*$ , then  $K = \text{span}(e_2, e_4, \dots, e_{2m})$  satisfies  $N \cap (K \otimes V^*) = 0$ .

For later use, we consider the space  $\Pi(\xi) \subset G(2, H^*)$  of all 2-dimensional subspaces  $K \subset H^*$  which contain  $\xi$ . We have:

**4.7. Lemma:** *Let  $n = 5$  and  $\omega \in M(5, 2)$  such that  $\mathcal{E}_\omega$  is not special 'tHooft. Then for a general  $\xi \in H^*$  there are closed subsets  $T_1 \subset T_2 \subset \Pi(\xi)$ ,  $\dim T_1 \leq 1$  and  $T_2$  a surface, such that*

- (i)  $N \cap (K \otimes V^*) = 0$  for  $K \in \Pi(\xi) \setminus T_2$ ;
- (ii)  $\dim N \cap (K \otimes V^*) = 1$  and  $N \cap (K \otimes V^*)$  contains no non-zero decomposable vector of  $H^* \otimes V^*$  for  $K \in T_2 \setminus T_1$ .

*Proof.* We use the previous notation and consider the morphisms

$$\mathbb{P}(N)_1 \xrightarrow{p_1} \mathbb{P}(H^*) \text{ and } \mathbb{P}(N)_2 \setminus \mathbb{P}(N)_1 \xrightarrow{p_2} G(2, H^*) =: G.$$

a) Let  $Y_1 \subset G$  denote the subset of those  $K \in G$  for which  $N \cap (K \otimes V^*)$  contains a non-zero decomposable vector or, equivalently,  $\mathbb{P}(K) \cap p_1(\mathbb{P}(N)_1) \neq \emptyset$ . Because  $\dim p_1(\mathbb{P}(N)_1) \leq 1$ , the set of lines  $\mathbb{P}(K)$  in  $\mathbb{P}(H^*)$  with this condition is closed and of dimension  $\leq 4$ .

b) Let  $Y_2 \subset G$  be the set of all  $K \in G$  for which  $N \cap (K \otimes V^*) \neq 0$ . If  $\mathcal{K} \subset H^* \otimes \mathcal{O}_G$  denotes the universal subbundle, we consider the homomorphism

$$\mathcal{K} \otimes V^* \xrightarrow{\phi} Q \otimes \mathcal{O}_G,$$

obtained as the composition of  $\mathcal{K} \otimes V^* \subset H^* \otimes V^* \otimes \mathcal{O}_G$  and the multiplication map  $H^* \otimes V^* \rightarrow Q$  with kernel  $N$ . Then  $Y_2$  is the determinantal subvariety of  $\phi$ . Because, for a general  $K$  we have  $N \cap (K \otimes V^*) = 0$  by Proposition 4.5,  $Y_2$  is a hypersurface in  $G$ ,  $\dim Y_2 = 5$ . By definition of  $Y_2$  we have  $\text{Im}(p_2) \subset Y_2$ , and, furthermore,

$$Y_2 = Y_1 \cup \text{Im}(p_2).$$

c) For  $K \in Y_2 \setminus Y_1$  the fibre  $p_2^{-1}(K)$  consists of vectors  $\xi_1 \otimes z_1 + \xi_2 \otimes z_2 \in N$ , where  $\xi_1, \xi_2$  is a basis of  $K$  and  $z_1, z_2 \in V^*$ , i.e.

$$p_2^{-1}(K) \cong \mathbb{P}(N \cap (K \otimes V^*)).$$

Now, by the proof of Proposition 4.5,  $p_2(\mathbb{P}(N)_2 \setminus \mathbb{P}(N)_1)$  has dimension  $\leq 5$ . It follows that, for a general  $K$  in each component of  $Y_2 \setminus Y_1$ , the fibre  $p_2^{-1}(K)$  is a point, or, equivalently,  $N \cap (K \otimes V^*)$  is 1-dimensional. Then the subvariety  $Y'_2 \subset Y_2$ , defined by  $\dim N \cap (K \otimes V^*) \geq 2$ , has dimension  $\leq 4$ . Let  $Z_1 = Y_1 \cup Y'_2$ . Then  $\dim Z_1 \leq 4$  and for  $K \in Y_2 \setminus Z_1$  the intersection  $N \cap (K \otimes V^*)$  is 1-dimensional and contains no non-zero decomposable vector.

d) Because  $\dim Z_1 \leq 4$ , the general 3-space  $\Pi(\xi)$  meets  $Z_1$  at most in dimension 1. Let, now,  $T_1 = \Pi(\xi) \cap Z_1$  and  $T_2 = \Pi(\xi) \cap Y_2$ . We may assume that  $\Pi(\xi) \not\subset Y_2$  by Proposition 4.5, b), so that  $T_2$  is a surface in  $\Pi(\xi)$ . Then  $T_1 \subset T_2$  satisfy the properties of the lemma.  $\square$

5. THE VARIETIES  $M(n, r)$  FOR  $n \leq 4$ 

We first note that  $M(n, 2n)$  is an open subset of the affine space  $S^2 H^* \otimes \Lambda^2 V^*$ . In fact, it is the complement of the hypersurface  $M_{4n-2}$ . In this case, if  $\omega \in M(n, 2n)$ , the bundle  $\mathcal{E}_\omega$  is the cokernel in

$$0 \rightarrow H \otimes \mathcal{O}_{\mathbb{P}}(-1) \rightarrow H^* \otimes \Omega^1(1) \rightarrow \mathcal{E}_\omega \rightarrow 0,$$

see 2.5 with  $Q = 0$ . Then  $H^1 \mathcal{E}_\omega(i) = 0$  for  $i \geq 0$  and we have

$$M^0(n, 2n) = M(n, 2n).$$

**5.1. Proposition:** *For  $n = 2$ ,  $M_6 = M_{4n-2}(H)$  is an irreducible hypersurface in  $S^2 H^* \otimes \Lambda^2 V^*$ .*

*Proof.* Note that  $M_6 \setminus (M_4 \cup \Delta)$  had been shown in 2.11 to be a smooth hypersurface, and  $M_6$  is a homogeneous hypersurface in  $S^2 H_2^* \otimes \Lambda^2 V^*$ .

In order to prove that it is irreducible, it suffices to prove that it is non-singular in codimension 1. Now  $\Delta = \Delta(H_2)$  is closed and irreducible in  $S^2 H_2^* \otimes \Lambda^2 V^*$  of codimension 2. We have  $M_4 \subset \Delta$  and hence  $M_6 \setminus \Delta$  is smooth. It is therefore sufficient to find an  $\omega \in \Delta \setminus M_4$  which is a smooth point of  $M_6$ . For that choose an isomorphism  $H_2 \cong k^2$  and let  $H_2 \xrightarrow{\omega} H_2^* \otimes \Lambda^2 V^*$  be presented by the matrix

$$\begin{pmatrix} x_2 \wedge x_3 & x_2 \wedge x_4 \\ x_2 \wedge x_4 & x_1 \wedge x_2 + x_3 \wedge x_4 \end{pmatrix}$$

where  $x_1, \dots, x_4$  is a basis of  $V^*$ . One can easily check that this  $\omega$  is degenerate at the point  $x_2 = x_3 = x_4 = 0$  and that  $\text{rk}(\omega) = 6$ . Moreover,  $\omega$  is a smooth point of  $M_6$ , using the argument of Katsylo–Ottaviani as in the proof of Proposition 2.11.  $\square$

**5.2. Corollary 1:** (Hartshorne)  *$MI(2)$  is smooth and irreducible of the expected dimension 13.*

*Proof.*  $M(2, 2) = M_6 \setminus (M_4 \cup \Delta)$  is a smooth transversal intersection and irreducible. By 2.9 the results follows.  $\square$

**5.3. Corollary 2:** (Ellingsrud - Strømme)  *$MI(3)$  is smooth and irreducible of the expected dimension 21.*

*Proof.* As noted above  $M^0(2, 4) = M(2, 4)$ , and this is irreducible. By Proposition 4.5  $M(3, 2)$  is the union of the open sets  $M(3, 2)_\xi$  which are equal to  $M(3, 2)'_\xi$  and which are transversal intersections and irreducible. The result follows now as in the previous case.  $\square$



In order to treat the case  $MI(4)$ , we prove the following

**5.4. Lemma:** *For  $n = 3$ ,  $M_{10} = M_{4n-2}(H)$  is an irreducible hypersurface in  $S^2H^* \otimes \Lambda^2V^*$ .*

*Proof.* By 2.11  $M_{10}$  is a hypersurface in  $S^2H_3^* \otimes \Lambda^2V^*$  and  $M(3, 4) = M_{10} \setminus (\Delta \cup M_8)$  is smooth. Because for  $n = 3$  we have  $M_6 \subset \Delta$ ,  $M(3, 2) = M_8 \setminus \Delta$  and hence  $\Delta \cup M_8 = \Delta \cup M(3, 2)$ . Now  $\Delta$  is irreducible in  $S^2H_3^* \otimes \Lambda^2V^*$  of codimension 4, see remark in 2.1, and  $M(3, 2)$  has codimension 6 in  $S^2H_3^* \otimes \Lambda^2V^*$  by corollary 5.3. It follows that  $M_{10}$  is smooth in codimension 1 and so  $M_{10}$  is irreducible.  $\square$

**5.5. Corollary:**  *$M(3, 4)$  is irreducible and a (smooth) transversal intersection of expected dimension 45.*

**5.6. Proposition:** (Barth, LePotier)  *$MI(4)$  is smooth and irreducible of the expected dimension 29.*

*Proof.* 1) Recall that the open subsets  $M(4, 2)'_\xi \subset M(4, 2)$  are defined by

$$M(4, 2)'_\xi = \{\omega \in M(4, 2) \mid \bar{\omega} \in M(3, 4) \text{ and } H^1\mathcal{E}_{\bar{\omega}}(1) = 0\},$$

where  $\bar{\omega} = \text{res}_\xi(\omega)$ . In this proof we write  $\mathcal{F} = \mathcal{E}_{\bar{\omega}}$ , which depends on the choice of  $\xi$ . By 3.3  $M(4, 2)'_\xi$  is irreducible. It is also smooth: by 2.8.1  $H^2S^2\mathcal{F} = 0$  for  $\bar{\omega} \in M(3, 4)$  because  $M(3, 4)$  is a smooth transversal intersection of the expected (co)dimension. It follows from Lemma 3.2 that also  $H^2S^2\mathcal{E}_\omega = 0$  for  $\omega \in M(4, 2)'_\xi$ , hence again by 2.8.1  $M(4, 2)'_\xi$  is smooth at any of its points and of expected dimension.

2) It suffices now to prove that

$$M(4, 2) = \bigcup_{\xi \neq 0} M(4, 2)'_\xi.$$

Let  $\omega \in M(4, 2)$ . By 6.3 there is  $\xi \in H^1\mathcal{E}_\omega(-1)$  such that  $H^1\mathcal{F}_P(1) = 0$  for any plane  $P$  in  $\mathbb{P}_3$ , provided  $\mathcal{E}_\omega$  is not special 'tHooft. In case  $n = 4$  we obtain  $h^1\mathcal{F} = 2$ . Then the operator  $H^1\mathcal{F} \otimes V^* \rightarrow H^1\mathcal{F}(1)$  can be presented by a matrix

$$A = \begin{pmatrix} v_{11} & \cdots & v_{1m} \\ v_{21} & \cdots & v_{2m} \end{pmatrix}$$

with  $m = h^1\mathcal{F}(1)$  if  $h^1\mathcal{F}(1) \neq 0$ . Let now  $P$  be a plane with equation  $z = 0$  and such that  $z(v_{11}) = z(v_{21}) = 0$ . Then

$$H^1\mathcal{F} \xrightarrow{A(z)} H^1\mathcal{F}(1)$$

cannot be surjective and it would follow that  $H^1\mathcal{F}_P(1) = \text{coker}A(z) \neq 0$ , a contradiction. This proves that  $\omega \in M(4, 2)'_\xi$  for the chosen  $\xi$ . If, however,  $\mathcal{E}_\omega$  is special 'tHooft, the pairing  $H^1\mathcal{E}_\omega(-1) \otimes V^* \rightarrow H^1\mathcal{E}_\omega$  can be presented by a  $4 \times 6$  matrix

$$\begin{pmatrix} v_1v_2 & & & & & \\ v_3v_4 & v_1v_2 & & & & \\ & v_3v_4 & v_1v_2 & & & \\ & & & v_3v_4 & & \end{pmatrix},$$

where  $v_1, \dots, v_4$  is a basis of  $V$ , see [6]. Then, choosing  $\xi = (0, 1, 0, 0) \in k^4 \cong H^1\mathcal{E}_\omega(-1)$ , the resulting homomorphism  $H^1\mathcal{F}(-1) \otimes V^* \rightarrow H^1\mathcal{F}$  is presented by the matrix

$$A = \begin{pmatrix} 0 & 0 \\ v_1 & v_2 \\ v_3 & v_4 \end{pmatrix}.$$

It follows that  $H^1\mathcal{F}(1) = 0$  because of the exact sequence  $H^1\mathcal{F}(-1) \otimes \Lambda^2 V^* \rightarrow H^1\mathcal{F} \otimes V^* \rightarrow H^1\mathcal{F}(1) \rightarrow 0$ .  $\square$

**5.7. Remark:** Concerning  $MI(5)$  and  $M(4, 4)$ , it follows from 3.3 that the open set

$$\bigcup_{\xi \neq 0} M(4, 4)_\xi$$

is irreducible and a (smooth) transversal intersection of expected dimension, because  $M^0(3, 6) = M(3, 6)$  is smooth of expected dimension. This will suffice to prove that  $MI(5)$  is smooth and irreducible of dimension 37. Using the method of Katsylo–Ottaviani as in the proof of 2.11, one can show that  $M(4, 4)$  is smooth of the expected dimension. It is an open question whether it is also irreducible.

## 6. A TECHNICAL RESULT ABOUT 5-INSTANTONS

**6.1. Proposition:** *Let  $\mathcal{E}_\omega$  be a 5-instanton on  $\mathbb{P}_3$ . Then for a general  $\xi \in H^1\mathcal{E}_\omega(-1)$  the associated rank-4 bundle  $\mathcal{F} = \mathcal{E}_{\bar{\omega}}$  satisfies  $h^1\mathcal{F}(1) \leq 1$ .*

For the proof we need the following lemmata on the vanishing of  $H^1\mathcal{F}_L(1)$  and  $H^1\mathcal{F}_P(1)$  for lines and planes.

**6.2. Lemma:** *Let  $\mathcal{E}_\omega$  be a  $n$ -instanton,  $n = 4$  or  $5$ , and assume that  $\mathcal{E}_\omega$  is not a special 'tHooft bundle ( $h^0\mathcal{E}_\omega(1) < 2$ ). Then, for a general  $\xi \in H^1\mathcal{E}_\omega(-1)$ , the bundle  $\mathcal{F} = \mathcal{E}_{\bar{\omega}}$  has the property that  $H^1\mathcal{F}_L(1) = 0$  for any line  $L$  in  $\mathbb{P}_3$ .*

*Proof.* Let  $\mathcal{E} = \mathcal{E}_\omega$ . For any plane  $P$  in  $\mathbb{P}_3$  we have  $H^1\mathcal{E}(-1) \xrightarrow{\sim} H^1\mathcal{E}_P(-1)$ . Then for any line  $L \subset P$  the restriction  $H^1\mathcal{E}(-1) \rightarrow H^1\mathcal{E}_L(-1)$  is surjective because  $H^1\mathcal{E}_P(-1) \rightarrow H^1\mathcal{E}_L(-1)$  is surjective, since  $H^1\mathcal{E}_P(-2) = 0$ . Because  $\mathcal{E}_L$  is the cohomology of the monad

$$0 \rightarrow \mathcal{O}_L(-1) \rightarrow \mathcal{F}_L \rightarrow \mathcal{O}_L(1) \rightarrow 0,$$

we obtain

$$H^1\mathcal{F}_L(1) \cong H^1\mathcal{E}_L(1)/\xi_L \cdot H^0\mathcal{O}_L(2),$$

where  $\xi_L$  denotes the restriction of  $\xi$  in  $H^1\mathcal{E}_L(-1)$ . On the other hand,  $\mathcal{E}_L \cong \mathcal{O}_L(a) \oplus \mathcal{O}_L(-a)$ ,  $0 \leq a \leq n$ . If  $0 \leq a \leq 2$ , then  $H^1\mathcal{E}_L(1) = 0$ . If  $a = 3$ ,  $H^1\mathcal{E}_L(-1) \cong H^1\mathcal{O}_L(-4)$  and  $H^1\mathcal{E}_L(1) \cong H^1\mathcal{O}_L(-2)$ . Then for  $\eta \in H^1\mathcal{O}_L(-4)$  we have  $\eta \cdot H^0\mathcal{O}_L(2) \neq H^1\mathcal{O}_L(-2)$  if and only if  $\eta = 0$ . Consequently, for the vanishing of  $H^1\mathcal{F}_L(1)$  we have only to assume that  $\xi_L \neq 0$  or equivalently that

$$\xi \notin \text{Ker}(H^1\mathcal{E}(-1) \rightarrow H^1\mathcal{E}_L(-1)),$$

which is 1- or 2-dimensional. Since by 4.3 the set of jumping lines of  $\mathcal{E}$  of order  $\geq 3$  is at most 1-dimensional,  $\xi$  has to avoid a subvariety of dimension  $\leq 2$  or 3. If  $a = 4$ , the elements  $\xi$  or  $\eta \in H^1\mathcal{O}_L(-5)$  should avoid the condition  $\eta H^0\mathcal{O}_L(2) \neq H^1\mathcal{O}_L(-3)$ . The set of these  $\eta$  is the affine cone over the rational normal curve in  $\mathbb{P}H^1\mathcal{O}_L(-5)$ . Namely, if  $s, t$  are homogeneous coordinates on  $L$ , we have

$$\eta = \sum_{\nu=1}^4 \frac{a_\nu}{s^{5-\nu}t^\nu} \quad \text{in} \quad H^1\mathcal{O}_L(-5)$$

and

$$s^2\eta = \frac{a_1}{s^2t} + \frac{a_2}{st^2}, \quad st\eta = \frac{a_2}{s^2t} + \frac{a_3}{st^3}, \quad t^2\eta = \frac{a_3}{s^2t} + \frac{a_4}{st^2}.$$

Then the condition  $\eta H^0\mathcal{O}_L(2) \neq H^1\mathcal{O}_L(-3)$  becomes

$$\text{rk} \begin{pmatrix} a_1 & a_2 & a_3 \\ a_2 & a_3 & a_4 \end{pmatrix} \leq 1.$$

So  $\xi$  has to avoid another 3-dimensional subvariety in  $H^1\mathcal{E}(-1)$ . Finally, if  $a = 5$ , in case  $n = 5$  only, the set of  $\eta \in H^1\mathcal{O}_L(-6)$  with  $\eta H^0\mathcal{O}_L(2) \neq H^1\mathcal{O}_L(-4)$  is a cubic hypersurface in  $H^1\mathcal{O}_L(-6)$ , as can be seen by a similar argument. So  $\xi$  has to avoid this hypersurface in  $H^1\mathcal{E}(-1) \cong H^1\mathcal{E}_L(-1)$  in case  $a = 5$ . But  $\mathcal{E}$  has only finitely many jumping lines of order 5 because it is not special 'tHooft, see also [24], [22]. Totally  $\xi$  has to avoid two 3-dimensional and finitely many 4-dimensional subvarieties in  $H^1\mathcal{E}(-1)$ , in case  $n = 5$ , in order to satisfy the condition of the lemma.  $\square$

**6.3. Lemma:** *Let  $\mathcal{E} = \mathcal{E}_\omega$  be as in the previous lemma. Then, for a general  $\xi \in H^1\mathcal{E}_\omega(-1)$ , the bundle  $\mathcal{F} = \mathcal{E}_\omega$  has the property that  $H^1\mathcal{F}_P(1) = 0$  for any plane  $P$  in  $\mathbb{P}_3$ .*

*Proof.* By 6.2 we may assume that  $H^1\mathcal{F}_L(1) = 0$  for any line  $L$  if  $\xi$  is in a fixed open set of  $H^1\mathcal{E}(-1)$ . We let  $\xi_P$  denote the element corresponding to  $\xi$  under the isomorphism  $H^1\mathcal{E}(-1) \cong H^1\mathcal{E}_P(-1)$  for a plane  $P$ . We are going to show that  $H^1\mathcal{F}_P(1) = 0$  for any plane if  $\xi$  avoids some additional subvarieties of  $H^1\mathcal{E}(-1)$ . These subvarieties will be estimated in dimension in the following cases.

**case 1:**  $P$  is a stable plane of  $\mathcal{E}$ . In this case there is no condition on  $\xi$  because we show that then already  $H^1\mathcal{F}_P(1) = 0$ .

Proof of case 1: We prove first that  $h^1\mathcal{F}_P \leq 2$ . As in the previous proof we have the exact sequence

$$\xi_P \otimes H^0\mathcal{O}_P(1) \xrightarrow{m(\xi)} H^1\mathcal{E}_P \rightarrow H^1\mathcal{F}_P \rightarrow 0.$$

Because  $h^0\mathcal{E}_P = 0$ , we have  $h^1\mathcal{E}_P = n - 2$ . So, if  $n = 4$ , then  $h^1\mathcal{F}_P \leq 2$ . If  $n = 5$ , the homomorphism  $m(\xi)$  is zero if and only if  $\xi_P$  is mapped to zero under the map  $H^1\mathcal{E}_P(-1) \rightarrow W \otimes H^1\mathcal{E}_P$ , where  $W^* = H^0\mathcal{O}_P(1)$ , and the kernel of this map is the image of  $\text{Hom}(\Omega_P^1(1), \mathcal{E}_P)$  in  $\text{Ext}^1(\mathcal{O}_P(1), \mathcal{E}_P) \cong H^1\mathcal{E}_P(-1)$ . Therefore,  $\xi_P$  is induced by a non-zero homomorphism  $\Omega_P^1(1) \xrightarrow{\varphi} \mathcal{E}_P$ . Because both bundles are stable,  $\varphi$  is generically and then globally injective. Then  $\mathcal{C} = \text{coker}(\varphi)$  has Hilbert polynomial  $\chi_{\mathcal{C}}(m) = m - 3$  and is Cohen–Macaulay. It follows that  $\mathcal{C} = \mathcal{O}_L(-4)$  for a line  $L \subset P$ . It follows that  $L$  is a jumping line of order 4 and that the restricted homomorphism  $\Omega_P^1 \otimes \mathcal{O}_L \rightarrow \mathcal{E}_L(-1)$  factors through  $\mathcal{O}_L(3)$ . Then the diagram

$$\begin{array}{ccc} H^1\Omega_P^1 & \longrightarrow & H^1\mathcal{E}_P(-1) \\ \downarrow & & \downarrow \\ H^1(\Omega_P^1 \otimes \mathcal{O}_L) & \longrightarrow & H^1\mathcal{E}_L(-1) \end{array}$$

implies that the image  $\xi_L$  of  $\xi$  in  $H^1\mathcal{E}_L(-1)$  is zero, and  $H^1\mathcal{F}_L(1) \cong H^1\mathcal{E}_L(1)/\xi_L H^0\mathcal{O}_L(2) \cong H^1\mathcal{O}_L(-3) \neq 0$  contradicting Lemma 6.2. This proves that  $h^1\mathcal{F}_P \leq 2$  if  $\xi$  is general.

Let now  $L \subset P$  be any line with equation  $z$ . By the assumption on  $\xi$  the multiplication map  $H^1\mathcal{F}_P \xrightarrow{z} H^1\mathcal{F}_P(1)$  is surjective. Applying the bilinear map lemma of H. Hopf to

$$H^1(\mathcal{F}_P(1))^* \otimes H^0\mathcal{O}_P(1) \rightarrow H^1(\mathcal{F}_P)^*,$$

we deduce that

$$h^1\mathcal{F}_P(1) \leq h^1\mathcal{F}_P - h^0\mathcal{O}_P(1) + 1 \leq 0.$$

(when  $K$  denotes the kernel of the map, the above surjectivity condition implies that  $\mathbb{P}H^1(\mathcal{F}(1))^* \times \mathbb{P}H^0\mathcal{O}_P(1)$  has an empty intersection with  $\mathbb{P}K$  in  $\mathbb{P}(H^1(\mathcal{F}(1))^* \otimes H^0\mathcal{O}_P(1))$ , which implies the estimate).

**case 2:**  $P$  is an unstable plane of  $\mathcal{E}$  but contains no jumping line of order 5.

In this case we have an exact sequence

$$0 \rightarrow \mathcal{O}_P \rightarrow \mathcal{E}_P \rightarrow \mathcal{I}_{Z,P} \rightarrow 0$$

where  $Z$  is a 0-dimensional subscheme of  $P$  of length  $n$  with  $H^0 \mathcal{I}_{Z,P}(1) = 0$ , because  $P$  contains no jumping line of order  $n$ . It follows that  $h^0 \mathcal{E}_P = 1$ ,  $h^0 \mathcal{E}_P(1) = 3$  and then  $h^1 \mathcal{E}_P = n-1$ ,  $h^1 \mathcal{E}_P(1) = n-3$ . If  $h^1 \mathcal{F}_P \leq 2$ , then, as in case 1,  $h^1 \mathcal{F}_P(1) = 0$ . If  $h^1 \mathcal{F}_P \geq 3$ , then the multiplication map  $m(\xi)$ , see case 1, has rank  $n-4$ . If  $n=4$ , then  $m(\xi) = 0$ , and we get a contradiction by the argument in case 1, which leads to  $h^1 \mathcal{F}_P \leq 2$ . If  $n=5$ , then  $m(\xi)$  has rank 1 and  $\xi_P$  is annihilated by two linear forms  $z_0, z_1 \in H^0 \mathcal{O}_P(1)$ . Let  $x \in P$  be the point determined by  $z_0, z_1$  and let  $\mathcal{I}_{\{x\},P}$  be its ideal sheaf. The standard resolution of this sheaf yields the exact sequence

$$0 \rightarrow H^0(\mathcal{E}_P)^2 \rightarrow H^0(\mathcal{I}_{\{x\},P} \otimes \mathcal{E}_P(1)) \rightarrow H^1 \mathcal{E}_P(-1) \xrightarrow{z_0, z_1} H^1(\mathcal{E}_P)^2.$$

On the other hand, the defining sequence of  $\mathcal{I}_{Z,P}$  and  $H^0 \mathcal{O}_P(1) \cong H^0 \mathcal{E}_P(1)$  implies

$$h^0(\mathcal{I}_{\{x\},P} \otimes \mathcal{E}_P(1)) = \begin{cases} 2 & \text{if } x \in P \setminus Z \\ 3 & \text{if } x \in Z \end{cases}.$$

Therefore,  $h^1 \mathcal{F}_P \geq 3$  can only occur if  $\xi_P$  avoids at most five 1-dimensional vector subspaces of  $H^1 \mathcal{E}_P(-1)$ . Because  $\mathcal{E}$  has only a 1-dimensional variety of unstable planes, it follows that  $h^1 \mathcal{F}_P \leq 2$  and  $h^1 \mathcal{F}_P(1) = 0$  for any unstable plane, if  $\xi$  avoids an at most 2-dimensional subvariety of  $H^1 \mathcal{E}(-1)$ .

**case 3:**  $P$  is an unstable plane of  $\mathcal{E}$  and contains a jumping line  $L$  of order  $n$ .

In this case  $H^1 \mathcal{E}_P(-1) \xrightarrow{\sim} H^1 \mathcal{E}_L(-1)$  and  $h^1 \mathcal{E}_P(1) = n-2 = h^1 \mathcal{E}_L(1)$ , such that  $H^1 \mathcal{E}_P(1) \xrightarrow{\sim} H^1 \mathcal{E}_L(1)$ . It follows that also  $H^1 \mathcal{F}_P(1) \xrightarrow{\sim} H^1 \mathcal{F}_L(1) = 0$ .  $\square$

**Proof of proposition 6.1:** Let us assume, firstly, that  $\mathcal{E} = \mathcal{E}_\omega$  is not special 'tHooft. By 4.5 we may assume that the multiplication map  $\xi \otimes V^* \rightarrow H^1 \mathcal{E}$  is injective, hence  $h^1 \mathcal{F} = 4$ ,  $\mathcal{F} = \mathcal{E}_\omega$ . By 6.3 we may assume that  $H^1 \mathcal{F}_P(1) = 0$  for any plane. Therefore, the multiplication map  $H^1 \mathcal{F} \rightarrow H^1 \mathcal{F}(1)$  is surjective for any linear form. Now an application of the bilinear lemma of H. Hopf to

$$H^1(\mathcal{F}(1))^* \otimes V^* \rightarrow H^1(\mathcal{F})^*$$

implies  $h^1(\mathcal{F}(1)) \leq h^1 \mathcal{F} - 4 + 1 = 1$ . If  $\mathcal{E}$  is a special 'tHooft bundle, it can even be shown that for a general  $\xi \in H^1 \mathcal{E}(-1)$  we have  $H^1 \mathcal{F}(1) = 0$ . If  $\mathcal{E}$  is special 'tHooft, then the pairing  $V^* \otimes H^1 \mathcal{E}(-1) \rightarrow H^1 \mathcal{E}$  can be defined by a  $5 \times 8$  matrix

$$\begin{pmatrix} v_1 & v_2 & & & & & & \\ v_3 & v_4 & v_1 & v_2 & & & & \\ & & v_3 & v_4 & v_1 & v_2 & & \\ & & & & v_3 & v_4 & v_1 & v_2 \\ & & & & & & v_3 & v_4 \end{pmatrix}$$

where  $v_1, \dots, v_4$  is a basis of  $V$ , after choosing suitable bases of the cohomology spaces, see [6]. Then, choosing  $\xi = (0, 0, 1, 0, 0)$  in  $k^5 \cong H^1\mathcal{E}(-1)$ , the resulting homomorphism  $V^* \otimes H^1\mathcal{F}(-1) \rightarrow H^1\mathcal{F}$  is described by the matrix

$$\left( \begin{array}{c|c} v_1v_2 & 0 \\ v_3v_4 & \\ \hline 0 & v_1v_2 \\ & v_3v_4 \end{array} \right)$$

which describes at the same time the right part  $4\Omega^1(1) \rightarrow 4\mathcal{O}$  of the Beilinson II monad of  $\mathcal{F}$ . It follows that  $h^1\mathcal{F}(1) = 0$  in this case.

## 7. IRREDUCIBILITY OF $MI(5)$

In this section the irreducibility of  $MI(5)$  is proved. The proof is mainly based on the properties of the sets  $M(4, 4)_\xi$  and the fact that for any  $\mathcal{E}_\omega \in MI(5)$  there is a  $\xi \in H^1\mathcal{E}_\omega(-1)$  such that  $(4, 4)$ -instanton  $\mathcal{E}_\omega$  satisfies  $h^1\mathcal{E}_\omega(1) \leq 1$ . The **plan of the proof** is the following. Let us recall the notations

$$M(4, 4) = \{\omega \in S^2H_4^* \otimes \Lambda^2V^* \mid \text{rk}(\omega) = 12, \omega \text{ non-degenerate}\}$$

and

$$M^0(4, 4) = \{\omega \in M(4, 4) \mid H^1\mathcal{E}_\omega(1) = 0\}.$$

In addition we need a partial completion of  $M(4, 4)$ , by forgetting the non-degeneracy,

$$\widetilde{M}(4, 4) := \{\omega \in S^2H_4^* \otimes \Lambda^2V^* \mid \text{rk}(\omega) = 12\}$$

and for  $\xi \in H_4^*$ , consistent with the earlier notation,

$$\widetilde{M}(4, 4)_\xi := \{\omega \in \widetilde{M}(4, 4) \mid \text{Im}(\omega) \cap (\xi \otimes V^*) = 0\}.$$

We have

$$\begin{aligned} M(4, 4)_\xi &= M(4, 4) \cap \widetilde{M}(4, 4)_\xi \\ M^0(4, 4)_\xi &= M^0(4, 4) \cap \widetilde{M}(4, 4)_\xi \end{aligned}$$

The space  $\widetilde{M}(4, 4)$  is introduced for technical reasons which become apparent in Lemma 7.1. For an element  $\omega \in \widetilde{M}(4, 4)$  we only get a sheaf  $\mathcal{E}_\omega$  from the monad construction 2.2. There is the morphism

$$\rho = \text{res}_\xi : \widetilde{M}(4, 4)_\xi \rightarrow M(3, 6) = M^0(3, 6)$$

assigning to  $\omega$  the map  $\bar{H}_3 \otimes V \xrightarrow{\bar{\omega}} \bar{H}_3^* \otimes V^*$ , where  $\bar{H}_3 = \text{Ker}(\xi)$ . Because  $\text{rk}(\bar{\omega}) = 12$ , this map is an isomorphism and hence non-degenerate. We are going to prove

**7.1. Lemma:**  $\widetilde{M}(4, 4)_\xi \xrightarrow{\rho} M(3, 6)$  is an affine bundle of dimension 54 and of fibre dimension 18.

Because  $M(3, 6)$  is an open part of  $S^2 \bar{H}_3^* \otimes \Lambda^2 V^*$ , it is smooth and irreducible of dimension 36, and therefore  $\widetilde{M}(4, 4)_\xi$  is smooth and irreducible of dimension 54. Then also the open subset  $M(4, 4)_\xi$  of  $\widetilde{M}(4, 4)_\xi$  is smooth and irreducible of dimension 54.

**Remark:** Irreducibility and smoothness of  $M(4, 4)_\xi$  is already proved 5.7. The bundle structure and the partial completion will be used to prove

**7.2. Lemma:** For any  $\xi \in H_4^* \setminus \{0\}$ ,  $\dim(M(4, 4)_\xi \setminus M^0(4, 4)_\xi) \leq \dim M(4, 4)_\xi - 2$ .

Assuming the two lemmata, the irreducibility of  $MI(5)$  is achieved with the following arguments. The open part

$$W := \bigcup_{\eta \neq 0} M(4, 4)_\eta$$

of  $M(4, 4)$  is smooth and irreducible by 3.4. We let

$$W^0 := \{\omega \in W \mid H^1 \mathcal{E}_\omega(1) = 0\} = \bigcup_{\eta \neq 0} M^0(4, 4)_\eta.$$

and

$$W^1 := \{\omega \in W \mid h^1 \mathcal{E}_\omega(1) \leq 1\}.$$

Then  $W^0 \subset W^1 \subset W$  are open subsets. It follows from Lemma 7.2, that

$$\text{codim}(W^1 \setminus W^0) \leq \dim W^1 - 2. \quad (1)$$

Now we consider the morphisms  $\omega \mapsto \bar{\omega} = \text{res}_\xi \omega$

$$M(5, 2)_\xi \rightarrow M(4, 4)$$

and the inverse images under  $r = \text{res}_\xi$ ,

$$\begin{array}{ccc} U_\xi^1 & \xrightarrow{r_1} & W^1 \\ \cup & & \cup \\ U_\xi^0 & \xrightarrow{r_0} & W^0. \end{array}$$

By Proposition 6.1, Lemma 3.1, and Proposition 4.5, any element  $\omega \in M(5, 2)$  is contained in one of the open sets  $U_\xi^1$ , i.e.

$$M(5, 2) = \bigcup_{\xi \neq 0} U_\xi^1,$$

but we don't know whether the same holds true for the sets  $U_\xi^0$ . By 3.3.1 the fibres of  $r_1$  and  $r_0$  are open sets of a linear space and their dimension are

$$\begin{aligned} \dim r_1^{-1}(\bar{\omega}) &= 4 + h^0 \mathcal{E}_{\bar{\omega}}(1) \leq 9 \\ \dim r_0^{-1}(\bar{\omega}) &= 4 + h^0 \mathcal{E}_{\bar{\omega}}(1) = 8. \end{aligned}$$

Because the difference of the fibre dimensions is at most 1, the estimate (1) implies

$$\dim(U_\xi^1 \setminus U_\xi^0) < \dim U_\xi^0 \quad (2)$$

for any  $\xi \neq 0$ . It follows that

$$\dim(M(5, 2) \setminus \bigcup_{\xi \neq 0} U_\xi^0) < \dim \bigcup_{\xi \neq 0} U_\xi^0 = 62.$$

Because any component of  $M(5, 2)$  has dimension  $\geq 25 + 40 - 3 = 62$  and because  $\bigcup_{\xi \neq 0} U_\xi^0$  is irreducible, it follows that  $M(5, 2)$  is irreducible of the expected dimension 62.

It follows now from Lemma 2.7.1 that also  $MI(5)$  is irreducible of the expected dimension 37. We thus have

**7.3. Theorem:**  *$MI(5)$  is irreducible of dimension 37.*

**7.4. Proof of Lemma 7.1:** We show in fact that

$$\widetilde{M}(4, 4)_\xi \cong M(3, 6) \times \text{Hom}_k(H_3, \Lambda^2 V^*).$$

For the proof we need to distinguish between the linear maps  $A \xrightarrow{\varphi} B \otimes \Lambda^2 V^*$  and the corresponding operators  $A \otimes V \xrightarrow{\tilde{\varphi}} B \otimes V^*$  which are skew with respect to  $V$  for any two vector spaces  $A$  and  $B$ . Because  $\xi$  is fixed, we can choose a decomposition

$$H_1 \oplus \bar{H}_3 = H_4,$$

where  $\bar{H}_3$  is the kernel of  $\xi$ . Given  $\bar{\omega} \in M(3, 6)$  and  $\bar{H}_3 \xrightarrow{\alpha} \Lambda^2 V^* \cong H_1^* \otimes \Lambda^2 V^*$ , we obtain the operator

$$\tilde{\omega} = \left( \begin{array}{c|c} \tilde{\alpha} \tilde{\omega}^{-1} \tilde{\alpha}^* & \tilde{\alpha} \\ \hline \tilde{\alpha}^* & \tilde{\omega} \end{array} \right) : (H_1 \oplus \bar{H}_3) \otimes V \rightarrow (H_1^* \oplus \bar{H}_3^*) \otimes V^*$$

where  $\alpha^*$  denotes the dual of  $\alpha$  with respect to  $\bar{H}_3$  and  $H_1$ . Because the upper row of  $\tilde{\omega}$  is a combination of the lower row, we have  $\text{rk}(\tilde{\omega}) = \text{rk}(\tilde{\omega}) = 12$ . It is clear that  $\tilde{\omega}$  is skew with respect to  $V$ . Therefore  $\tilde{\omega}$  defines an element  $\omega \in \widetilde{M}(4, 4)_\xi$ . We thus have a morphism

$$M(3, 6) \times \text{Hom}(\bar{H}_3, \Lambda^2 V^*) \rightarrow \widetilde{M}(4, 4)_\xi.$$

This is even an isomorphism over  $M(3, 6)$ , because if

$$\omega = \left( \begin{array}{c} \varphi \\ \alpha^* \end{array} \tilde{\omega} \right) : (H_1 \oplus \bar{H}_3) \rightarrow (H_1^* \oplus \bar{H}_3^*) \otimes \Lambda^2 V^*$$

is in  $M(4, 4)_\xi$ , we have  $12 = \text{rk}(\tilde{\omega}) = \text{rk}(\tilde{\omega})$  and it follows that

$$(\tilde{\varphi}, \tilde{\alpha}) = \tilde{\alpha} \tilde{\omega}^{-1} (\tilde{\alpha}^*, \tilde{\omega}).$$



**7.5. Proof of Lemma 7.2:** For fixed  $\eta \in H_4^* \setminus \{0\}$  there is an isomorphism

$$\widetilde{M}(4, 4)_\eta \cong M(3, 6) \times \text{Hom}(H_3, \Lambda^2 V^*) =: \widetilde{X}$$

by Lemma 7.1. To each pair  $(\bar{\omega}, \alpha) \in \widetilde{X}$  we have the simplified Beilinson II presentation of  $\mathcal{E}_{\bar{\omega}}$  together with a homomorphism  $\sigma(\alpha)$  induced by the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_3 \otimes \mathcal{O}(-1) & \xrightarrow{a(\bar{\omega})} & H_3^* \otimes \Omega^1(1) & \longrightarrow & \mathcal{E}_{\bar{\omega}} \longrightarrow 0 \\ & & \searrow \varepsilon & & \downarrow j & & \vdots \sigma(\alpha) \\ & & & & H_3^* \otimes V^* \otimes \mathcal{O} & & \vdots \\ & & & & \uparrow \bar{\omega} & & \vdots \\ & & & & H_3 \otimes V \otimes \mathcal{O} & \xrightarrow{\tilde{\alpha}} & \mathcal{O}(1) \end{array}$$

Note here that  $\tilde{\alpha} \circ \varepsilon = 0$  because  $\alpha$  is skew with respect to  $V$ , such that  $\tilde{\alpha} \circ \bar{\omega}^{-1} \circ j$  factors through  $\mathcal{E}_{\bar{\omega}}$ . Note further that in case  $(\bar{\omega}, \alpha)$  corresponds to an  $\omega \in M(4, 4)_\eta$ , then  $\mathcal{E}_\omega$  is the cohomology of the monad

$$\mathcal{O}(-1) \xrightarrow{\sigma(\alpha)^*} \mathcal{E}_{\bar{\omega}}^* \xrightarrow[j_{\bar{\omega}}]{\approx} \mathcal{E}_{\bar{\omega}} \xrightarrow{\sigma(\alpha)} \mathcal{O}(1)$$

in which case  $\sigma(\alpha)$  is surjective, see beginning of Section 3.

We let  $X \subset \widetilde{X}$  be the open subset corresponding to  $M(4, 4)_\eta$ , or defined by the surjectivity of  $\sigma(\alpha)$ . In addition we let  $\widetilde{X}^0 \subset \widetilde{X}$  denote the open part of  $\widetilde{X}$  where

$$H^0 \sigma(\alpha)(1) : H^0 \mathcal{E}_{\bar{\omega}}(1) \rightarrow H^0 \mathcal{O}(2)$$

is surjective. Then

$$M^0(4, 4)_\eta \cong X^0 = X \cap \widetilde{X}^0$$

under the above isomorphism because the cokernel of  $H^0 \sigma(\alpha)(1)$  is then isomorphic to  $H^1 \mathcal{E}_\omega(1)$ . To prove Lemma 7.2, it is sufficient to prove that

$$\text{codim}(\widetilde{X} \setminus \widetilde{X}^0, \widetilde{X}) \geq 2. \quad (*)$$

Now  $(*)$  will follow from the following two statements.

**Claim F:** There exists an  $\bar{\omega} \in M(3, 6)$  such that for the fibres,  $\widetilde{X}_{\bar{\omega}}^0 \subset \widetilde{X}_{\bar{\omega}}$  of  $\widetilde{X}^0 \subset \widetilde{X}$  over  $\bar{\omega}$ ,

$$\text{codim}(\widetilde{X}_{\bar{\omega}} \setminus \widetilde{X}_{\bar{\omega}}^0, \widetilde{X}_{\bar{\omega}}) \geq 2.$$

**Claim B:** Let  $\Sigma \subset M(3, 6)$  be the closed subvariety of points  $\bar{\omega}$  for which  $\widetilde{X}_{\bar{\omega}}^0 = \emptyset$ , i.e. the set of points  $\bar{\omega}$  for which  $H^0 \sigma(\alpha)(1)$  is not surjective for any  $\alpha$ . Then

$$\text{codim}(\Sigma, M(3, 6)) \geq 2.$$

### 7.5.1. Proof of claim B:

a) In order to incorporate a projective curve which doesn't meet  $\Sigma$ , we enlarge  $M(3, 6)$  and  $\Sigma$  as follows. Let

$$\widetilde{M}(3, 6) := M(3, 6) \cup M^0(3, 4).$$

Then  $\widetilde{M}(3, 6) \subset S^2 H_3^* \otimes \Lambda^2 V^*$  consists of non-degenerate  $\bar{\omega}$  of rank 10 if  $\bar{\omega} \in M^0(3, 4)$  and of rank 12 if  $\bar{\omega} \in M(3, 6)$ , and such that  $H^1 \mathcal{E}_{\bar{\omega}}(1) = 0$ . If  $\bar{\omega} \in M^0(3, 4)$ , the Beilinson II monad of  $\mathcal{E}_{\bar{\omega}}$  is of the type

$$0 \rightarrow H_3 \otimes \mathcal{O}(-1) \xrightarrow{a(\bar{\omega})} H_3^* \otimes \Omega^1(1) \rightarrow Q \otimes \mathcal{O} \rightarrow 0$$

where  $Q$  is the cokernel of  $\bar{\omega}$ .

If  $\mathcal{F}_{\bar{\omega}}$  is the cokernel of  $a(\bar{\omega})$ , we have the exact sequence

$$0 \rightarrow \mathcal{E}_{\bar{\omega}} \rightarrow \mathcal{F}_{\bar{\omega}} \rightarrow Q \otimes \mathcal{O} \rightarrow 0.$$

Given a second component  $\alpha \in \text{Hom}(H_3, \Lambda^2 V^*)$ , we also obtain a homomorphism  $\mathcal{F}_{\bar{\omega}} \xrightarrow{\sigma(\alpha)} \mathcal{O}(1)$  by the diagram

$$\begin{array}{ccccccc} & & H_3 \otimes V \otimes \mathcal{O} & \xrightarrow{\tilde{\alpha}} & \mathcal{O}(1) & & \\ & & \downarrow \bar{\omega} & \nearrow & \uparrow & & \\ & & H_3^* \otimes V \otimes \mathcal{O} & & & & \\ & & \uparrow & & \uparrow \sigma(\alpha) & & \\ 0 & \longrightarrow & H_3 \otimes \mathcal{O}(-1) & \longrightarrow & H_3^* \otimes \Omega^1(1) & \longrightarrow & \mathcal{F}_{\bar{\omega}} \longrightarrow 0 \end{array}$$

which in case  $\bar{\omega} \in M(3, 6)$  coincides with  $\mathcal{E}_{\bar{\omega}} \xrightarrow{\sigma(\alpha)} \mathcal{O}(1)$ . The surjectivity of  $H^0 \sigma(\alpha)(1)$  does not depend on the choice of the factorization of  $\tilde{\alpha}$ .

We let now  $\widetilde{\Sigma} \subset \widetilde{M}(3, 6)$  be the locus of points  $\bar{\omega}$  for which  $H^0 \sigma(\alpha)(1)$  is not surjective for any  $\alpha$ . By this definition we have

$$\Sigma = \widetilde{\Sigma} \cap M(3, 6).$$

b) Because  $M_{10}(H_3)$  had been shown to be an irreducible hypersurface in  $S^2 H_3^* \otimes \Lambda^2 V^*$  whose complement is  $M(3, 6)$ , see 5.4, and because  $M^0(3, 4)$  is an open part of it, the complement of  $\widetilde{M}(3, 6)$  in  $S^2 H_3^* \otimes \Lambda^2 V^*$  has codimension  $\geq 2$ . In order to show that  $\text{codim}(\widetilde{\Sigma}, \widetilde{M}(3, 6)) \geq 2$ , we first construct an embedding  $k^2 \setminus \{0\} \rightarrow \widetilde{M}(3, 6) \setminus \widetilde{\Sigma}$ .

c) This embedding is defined as follows. We let  $e_1, \dots, e_4$  be the standard basis of  $H_4 = k^4$  and  $e_1^*, \dots, e_4^*$  be its dual basis. We let

$$H_4 \xrightarrow{\omega} H_4^* \otimes \Lambda^2 V^*$$

be given by the matrix

$$\omega = \begin{pmatrix} \omega'_{11} & \omega'_{12} & 0 & 0 \\ \omega'_{12} & \omega'_{22} & 0 & 0 \\ 0 & 0 & \omega''_{11} & \omega''_{12} \\ 0 & 0 & \omega''_{12} & \omega''_{22} \end{pmatrix}$$

with  $\omega'_{ij}, \omega''_{ij} \in \Lambda^2 V^*$  which represents the direct sum of two 2-instantons  $\mathcal{E}'$  and  $\mathcal{E}''$ ,  $\mathcal{E}_\omega = \mathcal{E}' \oplus \mathcal{E}''$ . Then  $H^1 \mathcal{E}_\omega(1) = 0$ . For  $t = (t_0, t_1) \neq 0$  we consider

$$\xi_t = (-t_1, 0, t_0, 0) \in H_4^*$$

and its kernel

$$0 \rightarrow k^3 \xrightarrow{f_t} k^4 \xrightarrow{\xi_t} k \rightarrow 0$$

defined by the matrix

$$f_t = \begin{pmatrix} t_0 & 0 & 0 \\ 0 & 1 & 0 \\ t_1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We let  $\bar{\omega}_t = (f_t^* \otimes \text{id}) \circ \omega \circ f_t$  such that

$$\bar{\omega}_t = \begin{pmatrix} t_0^2 \omega'_{11} + t_1^2 \omega''_{11} & t_0 \omega'_{12} & t_1 \omega''_{12} \\ t_0 \omega'_{12} & \omega'_{22} & 0 \\ t_1 \omega''_{12} & 0 & \omega''_{22} \end{pmatrix}.$$

We may assume that both  $\omega'_{22}$  and  $\omega''_{22}$  have rank 4 as operators  $V \rightarrow V^*$ . Then by an elementary matrix operation we can kill  $t_0 \omega'_{12}$  and  $t_1 \omega''_{12}$  in the first row and obtain a matrix

$$\begin{pmatrix} t_0^2 \eta'_{11} + t_1^2 \eta''_{11} & 0 & 0 \\ t_0 \omega'_{12} & \omega'_{22} & \\ t_1 \omega''_{12} & & \omega''_{22} \end{pmatrix}$$

with  $\eta'_{11} = \omega'_{11} - \omega'_{12} \omega'^{-1}_{22} \omega'_{12}$  and  $\eta''_{11} = \omega''_{11} - \omega''_{12} \omega''^{-1}_{22} \omega''_{12}$ .

Because  $\omega'$  and  $\omega''$  represent 2-instantons,  $\text{rk}(\omega') = \text{rk}(\omega'') = 6$  and therefore

$$\text{rk } \eta'_{11} = \text{rk } \eta''_{11} = 2.$$

But we can choose  $\omega'$  and  $\omega''$  such that  $\text{rk } \omega'_{22} = \text{rk } \omega''_{22} = 4$  and in addition

$$\text{Im } \eta'_{11} + \text{Im } \eta''_{11} = V^*.$$

Then  $t_0^2 \eta'_{11} + t_1^2 \eta''_{11}$  is an isomorphism  $V \rightarrow V^*$  for  $t_0 t_1 \neq 0$ . Therefore, with this choice of  $\omega'$  and  $\omega''$ , we have

$$\begin{aligned} \bar{\omega}_t &\in M(3, 6) && \text{for } t_0 t_1 \neq 0 \\ \bar{\omega}_t &\in M^0(3, 4) && \text{if } t_0 = 0 \text{ or } t_1 = 0. \end{aligned}$$

The first statement follows directly from the fact that  $\text{rk } \bar{\omega}_t = 12$  if  $t_0 t_1 \neq 0$ . If  $t_0 = 0$  or  $t_1 = 0$ , then  $\text{rk } \bar{\omega}_t = 10$ . In that case, e.g.  $t_1 = 0$ ,

$$\bar{\omega}_t = \begin{pmatrix} t_0^2 \omega'_{11} & t_0 \omega'_{12} & 0 \\ t_0 \omega'_{12} & \omega'_{22} & 0 \\ 0 & 0 & \omega''_{22} \end{pmatrix}$$

and then the bundle of  $\bar{\omega}_t$  is the direct sum

$$\mathcal{E}_{\bar{\omega}_t} \cong \mathcal{E}_{\omega'} \oplus \mathcal{E}_{\omega''_{12}}$$

with  $\mathcal{E}' = \mathcal{E}_{\omega'}$  a 2-instanton and  $\mathcal{E}_{\omega''_{22}}$  a 1-instanton (null-correlation bundle). So  $\bar{\omega}_t$  is non-degenerate and  $H^1 \mathcal{E}_{\bar{\omega}_t}(1) = 0$ . It follows that  $t \mapsto \bar{\omega}_t$  is a morphism

$$k^2 \setminus \{0\} \rightarrow \widetilde{M}(3, 6)$$

which is an embedding by the shape of the matrix  $\bar{\omega}_t$ . Moreover,  $\bar{\omega}_t \notin \widetilde{\Sigma}$  for any  $t$ . To see this, we consider the case  $t_1 \neq 0$  first. In that case the linear embedding  $k^3 \xrightarrow{f_t} k^4$  can be considered as the kernel of a fixed  $\xi = (1, 0, 0, 0)$  by the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & k^3 & \xrightarrow{f_t} & k^4 & \xrightarrow{\xi_t} & k \longrightarrow 0 \\ & & \parallel & & \downarrow \approx a_t & & \parallel \\ 0 & \longrightarrow & k^3 & \xrightarrow{g} & k^4 & \xrightarrow{\xi} & k \longrightarrow 0 \end{array}$$

where  $\xi_t = (-t_1, 0, t_0, 0)$  and  $\xi = (1, 0, 0, 0)$  with

$$g = \begin{pmatrix} 0 & 0 & 0 \\ 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} \quad \text{and} \quad a_t = \begin{pmatrix} -1 & 0 & t_0/t_1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

If we denote

$$\omega_t := a_t^{-1,*} \omega a_t^{-1}$$

we have

$$\bar{\omega}_t = g^* \omega_t g = \text{res}_\xi \omega_t.$$

Because  $\omega_t \sim \omega$ , we have  $\mathcal{E}_{\omega_t} \cong \mathcal{E}_\omega$  and so  $H^1 \mathcal{E}_{\omega_t}(1) = 0$ . If  $\bar{\omega}_t \in M(3, 6)$ , then  $\bar{\omega}_t \notin \Sigma$ . If, however,  $\bar{\omega}_t \in M^0(3, 4)$ , then  $\mathcal{E}_{\bar{\omega}_t}$  is the cohomology of the complex

$$\mathcal{O}(-1) \longrightarrow \mathcal{F}_{\bar{\omega}_t}^* \longrightarrow \mathcal{F}_{\bar{\omega}_t} \longrightarrow \mathcal{O}(1).$$

Then also in this case  $H^0 \mathcal{F}_{\bar{\omega}_t}(1) \rightarrow H^0 \mathcal{O}(2)$  has cokernel  $H^1 \mathcal{E}_{\omega_t}(1) = 0$ . Therefore  $\bar{\omega}_t \notin \widetilde{\Sigma}$  for  $t_1 \neq 0$ . By symmetry in  $t_0, t_1$ , also  $\bar{\omega}_t \notin \widetilde{\Sigma}$  for  $t_0 \neq 0$ . Therefore we have an embedding

$$k^2 \setminus \{0\} \hookrightarrow \widetilde{M}(3, 6) \setminus \widetilde{\Sigma}.$$

d) The map  $t \mapsto \bar{\omega}_t$  cannot directly be used to define a line in  $\mathbb{P}S^2H_3^* \otimes \Lambda^2V^*$  because  $\bar{\omega}_t$  is not homogeneous in  $t$ . But it can be used to construct a projective line in  $\mathbb{P}H_4^*$  which doesn't intersect a transformation of  $\tilde{\Sigma}$  so that we can conclude from that that  $\tilde{\Sigma}$  has codimension  $\geq 2$ . To do this we consider the following transformation diagram

$$\mathbb{P}H_4^* \xleftarrow{\pi} \mathbb{P}\mathrm{Hom}(H_3, H_4)^0 \xrightarrow{\theta} \mathbb{P}S^2H_3^* \otimes \Lambda^2V^*.$$

In this diagram  $\mathbb{P}\mathrm{Hom}(H_3, H_4)^0$  denotes that open set of injective maps and  $\pi$  is the principal  $PGL(H_3)$ -bundle over the Grassmannian  $G(3, H_4) = \mathbb{P}H_4^*$ . The morphism  $\theta$  is defined by

$$f \mapsto (f^* \otimes \mathrm{id}) \circ \omega \circ (f \otimes \mathrm{id})$$

using the form of  $\omega$  in c). We let

$$\widehat{\Sigma} \subset \widehat{M}(3, 6) \subset \mathbb{P}S^2H_3^* \otimes \Lambda^2V^*$$

denote the subvarieties of which  $\tilde{\Sigma}$  and  $\tilde{M}(3, 4)$  are the affine cones. So  $\widehat{\Sigma}$  is closed in the open set  $\widehat{M}(3, 6)$  of codimension  $\geq 2$ . Now

$$\theta^{-1}\widehat{\Sigma} \subset \theta^{-1}\widehat{M}(3, 6) \subset \mathbb{P}\mathrm{Hom}(H_3, H_4)^0$$

are  $PGL(H_3)$ -invariant and therefore there are an open subset  $U \subset \mathbb{P}H_4^*$  and a closed subscheme  $Z \subset U$  such that

$$\theta^{-1}\widehat{\Sigma} = \pi^{-1}Z \quad \text{and} \quad \theta^{-1}\widehat{M}(3, 6) = \pi^{-1}U.$$

Moreover, there is the irreducible hypersurface  $\widehat{M}_{10}(H_3) \subset \mathbb{P}S^2H_3^* \otimes \Lambda^2V^*$  defined by  $\mathrm{rk} \bar{\omega} \leq 10$ . Its inverse image  $\theta^{-1}\widehat{M}_{10}(H_3)$  is the subvariety of those  $\langle f \rangle \in \mathbb{P}\mathrm{Hom}(H_3, H_4)^0$ , for which  $(H_4/\mathrm{Im}(f))^* \otimes V^* \rightarrow H^1\mathcal{E}_\omega$  is not an isomorphism, see 3.1, because then  $\bar{\omega} = \theta(f)$  has rank 10. It is then easily seen that this condition defines an effective divisor, which is  $\theta^{-1}\widehat{M}_{10}(H_3)$ . Because this is also  $PGL(H_3)$ -invariant, there is a divisor  $D \subset \mathbb{P}H_4^*$  such that

$$\theta^{-1}\widehat{M}_{10}(H_3) = \pi^{-1}D.$$

Now the family  $(f_t)$  of c) defines a linear embedding  $\langle t \rangle \mapsto \pi(\langle f_t \rangle) = \langle \xi_t \rangle$

$$\mathbb{P}_1 \xrightarrow[\approx]{} L \subset \mathbb{P}H_4^*$$

and by the result of c) we have

$$L \subset U \setminus Z \quad \text{and} \quad L \not\subset D.$$

Let  $Y \subset \mathbb{P}H_4^*$  be the complement of  $U$ . Then  $L$  doesn't intersect  $Y \cup Z = Y \cup \bar{Z}$  and so  $Z$  is of codimension  $\geq 2$ . Then also  $\theta^{-1}\widehat{\Sigma}$  has codimension  $\geq 2$  in  $\theta^{-1}\widehat{M}(3, 6)$ . But since also the complement of  $\theta^{-1}\widehat{M}(3, 6)$  has codimension  $\geq 2$  in  $\mathbb{P}\mathrm{Hom}(H_3, H_4)$ , there is a line  $L' \subset \mathbb{P}\mathrm{Hom}(H_3, H_4)$  such that

$$L' \subset \theta^{-1}\widehat{M}(3, 6) \setminus \theta^{-1}\widehat{M}_{10}(H_3) \quad \text{and} \quad L' \not\subset \theta^{-1}\widehat{M}_{10}(H_3).$$

We let now  $\Gamma = \theta(L')$ . This is a complete curve in  $\mathbb{P}S^2H_3^* \otimes S^2V^*$  (not contracted to a point because  $L'$  intersects the divisor  $\theta^{-1}\widehat{M}_{10}(H_3)$  and is not contained in it) and such that

$$\Gamma \subset \widehat{M}(3, 6) \setminus \widehat{\Sigma}.$$

Then  $\Gamma$  cannot intersect the closure of  $\widehat{\Sigma}$  in  $\mathbb{P}S^2H_3^* \otimes \Lambda^2V^*$ . So we have shown that  $\text{codim}(\widehat{\Sigma}, \widehat{M}(3, 6)) \geq 2$ . This is equivalent to claim B.

### 7.5.2. Remarks on null-correlation bundles:

The tensors  $\bar{\omega}$  provided for claim F are diagonal matrices of elements  $\eta \in \Lambda^2V^*$  of rank 4 (or indecomposable) which correspond to 1-instantons or null-correlation bundles (nc-bundles for short). Given  $\eta$  of rank 4, we have the defining sequence  $0 \rightarrow \mathcal{O}(-1) \rightarrow \Omega^1(1) \rightarrow \mathcal{E}_\eta \rightarrow 0$ . Then for any  $\alpha \in \Lambda^2V^*$  we obtain a homomorphism  $\mathcal{E}_\eta \xrightarrow{\sigma(\alpha)} \mathcal{O}(1)$  as in 7.5.1, using the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}(-1) & \xrightarrow{\eta} & \Omega^1(1) & \longrightarrow & \mathcal{E}_\eta \longrightarrow 0 \\ & & & & \downarrow & & \vdots \\ & & & & V^* \otimes \mathcal{O} & & \vdots \\ & & & & \uparrow \eta & & \vdots \\ & & & & V \otimes \mathcal{O} & \xrightarrow{\tilde{\alpha}} & \mathcal{O}(1) \end{array}$$

One can easily see that  $\sigma(\alpha) = 0$  if and only if  $\alpha$  and  $\eta$  are proportional in  $\Lambda^2V^*$  or  $\langle \alpha \rangle = \langle \eta \rangle$ . Therefore, a non-zero image  $\mathcal{I}$  of  $\mathcal{E}_\eta(-1) \rightarrow \mathcal{O}$  is determined by a line  $g = \langle \alpha \rangle, \langle \eta \rangle$  in  $\mathbb{P}\Lambda^2V^*$  and is independent of  $\langle \alpha \rangle$  on  $g$ . We therefore write  $\mathcal{I}_g$  for the image of  $\mathcal{E}_\eta(-1)$  under  $\sigma(\alpha)$ .

In this section we identify the Grassmannians  $G = G(2, V)$  and  $G(2, V^*)$  via the Plücker embeddings in  $\mathbb{P}\Lambda^2V \cong \mathbb{P}\Lambda^2V^*$ , because  $\Lambda^2V \cong \Lambda^2V^* \otimes \Lambda^4V$  canonically. Then for a line  $l$  in  $\mathbb{P}V$  through  $\langle x \rangle, \langle y \rangle$  with equations  $z, w$  we write  $l = \langle x \wedge y \rangle = \langle z \wedge w \rangle$ .

**Lemma 1:** *Let  $\langle \eta \rangle \in \mathbb{P}\Lambda^2V^* \setminus G$  and let  $g$  be a line in  $\mathbb{P}\Lambda^2V^*$  through  $\langle \eta \rangle$  with associated ideal sheaf  $\mathcal{I}_g$ . Then*

- a) *If  $g$  meets the Grassmannian  $G$  in two points  $l_1 \neq l_2$ , then  $l_1 \cap l_2 = \emptyset$  in  $\mathbb{P}V$  and  $\mathcal{I}_g = \mathcal{I}_{l_1 \cup l_2}$ .*
- b) *If  $g$  is tangent to  $G$  at  $l$ , then  $\mathcal{I}_g$  is the twisted double structure on the line  $l$  given by the tangent direction along  $g$ .*

*In both cases  $h^0\mathcal{I}_g(2) = 4$ .*

*Proof.* Let  $Z$  be the zero scheme of  $\mathcal{I}_g$ . We have the exact sequence  $0 \rightarrow \mathcal{O}(-1) \rightarrow \mathcal{E}_\eta \rightarrow \mathcal{O}(1) \rightarrow \mathcal{O}_Z(1) \rightarrow 0$  and by that for the Hilbert polynomial  $\chi\mathcal{O}_Z(m) = 2m + 2$ .

This means that  $Z$  is a pair of disjoint lines or a twisted double structure on a line. In order to prove that  $\mathcal{I}_g$  is the ideal of the lemma, it is sufficient to determine the space  $H^0\mathcal{I}_g(2) \subset S^2V^*$  by the induced diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & k & \xrightarrow{\eta} & \Lambda^2 V^* & \longrightarrow & H^0\mathcal{E}_\eta(1) \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow H^0\sigma(\alpha)(1) \\
 & & & & V^* \otimes V^* & & \\
 & & & \approx \uparrow \eta \otimes \text{id} & & & \\
 & & V \otimes V^* & \xrightarrow{\tilde{\alpha}} & S^2 V^* & & 
 \end{array}$$

where  $g \ni \langle \alpha \rangle \neq \langle \eta \rangle$ .

**case a)** Because  $\langle \alpha \rangle, \langle \eta \rangle, l_1, l_2$  are on the same line  $g \not\subset G$ , we may choose a basis  $e_0, \dots, e_3 \in V$  with dual basis  $z_0, \dots, z_3$  such that  $l_1 = \langle e_0 \wedge e_1 \rangle$ ,  $l_2 = \langle e_2 \wedge e_3 \rangle$ ,  $\eta = z_0 \wedge z_1 + z_2 \wedge z_3$  and  $\alpha = \lambda z_0 \wedge z_1 + \mu z_2 \wedge z_3$  with  $\lambda \neq \mu$ . With this choice one computes that in the above diagram the image of  $H^0\sigma(\alpha)(1)$  is spanned by the 4 forms  $z_0 z_2, z_0 z_3, z_1 z_2, z_1 z_3$  which are the generators of the ideal of  $l_1 \cup l_2$ .

**case b)** Here we may choose the same  $\eta$  but  $\alpha = z_1 \wedge z_3$  such that  $\langle \alpha \rangle = \langle e_0 \wedge e_2 \rangle = l$  in  $\mathbb{P}\Lambda^2 V^* = \mathbb{P}\Lambda^2 V$ . With the same type of calculation we find that  $H^0\mathcal{I}_g(2)$  is spanned by the 4 forms  $z_1^2, z_1 z_3, z_3^2, z_0 z_1 + z_2 z_3$ , which are the generators of the ideal of the double structure on  $l$ .  $\square$

**Lemma 2:** (1) Let  $p_1, p_2$  be two different points on a line  $l$  in  $\mathbb{P}_3$  and let  $l_1, l_2$  be two skew lines not meeting  $p_1, p_2$ . Then the space of quadrics containing  $p_1, p_2, l_1, l_2$  is 2-dimensional if and only if  $l$  doesn't meet both  $l_1$  and  $l_2$ , and otherwise this space is 3-dimensional.

(2) If  $l_0$  is a twisted double line as in Lemma 1 above, which doesn't meet  $l$ , then the space of quadrics containing  $p_1, p_2$  and  $l_0$  is 2-dimensional, too.

*Proof.* By elementary computation using suitable coordinates.  $\square$

### 7.5.3. Three nc-bundles:

We consider first two nc-bundles  $\mathcal{E}_1 = \mathcal{E}_{\eta_1}, \mathcal{E}_2 = \mathcal{E}_{\eta_2}$  with homomorphisms  $\mathcal{E}_i \xrightarrow{\sigma_i} \mathcal{O}(1)$ , and ideal sheaves  $\mathcal{I}_1, \mathcal{I}_2$  respectively. Then the image of  $H^0(\mathcal{E}_1(1) \oplus \mathcal{E}_2(1)) \rightarrow H^0\mathcal{O}(2)$  under  $\sigma_1 + \sigma_2$  is  $H^0\mathcal{I}_1(2) + H^0\mathcal{I}_2(2)$ , and similarly for three such data  $(\mathcal{I}_i, \sigma_i)$ . We let  $Z_1, Z_2$  be the zero schemes of  $\mathcal{I}_1, \mathcal{I}_2$  respectively.

**Lemma 3:** Let  $Z_1 = l_{11} \cup l_{12}$  and  $Z_2 = l_{21} \cup l_{22}$  with all 4 lines different. Suppose that either

(i)  $Z_1 \cap Z_2 = \emptyset$  and the 4 lines do not belong to the same ruling of a quadric, or

(ii)  $Z_1 \cap Z_2 = \{x_1, x_2\} \subset l_{11}$  and  $Z_2 \cap l_{12} = \emptyset$ .

Then

$$H^0\mathcal{I}_1(2) \cap H^0\mathcal{I}_2(2) = 0.$$

*Proof.* (i) well-known. (ii) Let  $Q$  be a smooth quadric containing  $l_{11}, l_{12}$  (in one ruling). Because  $l_{2\nu} \cap l_{12} = \emptyset$  and  $l_{2\nu} \cap l_{11} = \{x_\nu\}$ , we have  $l_{21}, l_{22} \not\subset Q$ . Therefore the intersection of the  $H^0\mathcal{I}_\nu(2)$  does not contain a smooth quadric. Then any  $Q \in H^0\mathcal{I}_1(2) \cap H^0\mathcal{I}_2(2)$  is not smooth and contains the skew lines  $l_{11}$  and  $l_{12}$ . So  $Q = P_1 \cup P_2$  is a pair of planes. Let  $l_{11} \subset P_1$ . Then  $l_{12} \not\subset P_1$  because  $l_{11} \cap l_{12} = \emptyset$  and so  $l_{12} \subset P_2$ . Then  $l_{21}, l_{22} \not\subset P_2$  and  $l_{21}, l_{22} \subset P_1$ . But  $l_{21} \cap l_{22} = \emptyset$ , a contradiction.  $\square$

**Remark:** In case (ii) the 8-dimensional space  $H^0\mathcal{I}_1(2) + H^0\mathcal{I}_2(2)$  equals  $H^0\mathcal{I}_{\{x_1, x_2\}}(2)$ .

**Corollary:** Let the situation be as in Lemma 3 and let  $\mathcal{E}_3 \xrightarrow{\sigma_3} \mathcal{O}(1)$  be a third homomorphism of an nc-bundle with ideal  $\mathcal{I}_3$  and zero scheme  $Z_3$ . If  $Z_3$  consists of two (then skew) lines  $l_{31}, l_{32}$  or of a twisted double line  $l_3$ , such that the line

$$l_{11} = \overline{x_1, x_2}$$

doesn't meet both of  $l_{31}$  and  $l_{32}$ , then

$$H^0\mathcal{I}_1(2) + H^0\mathcal{I}_2(2) + H^0\mathcal{I}_3(2) = H^0\mathcal{O}(2).$$

*Proof.* By Lemma 3 the sum of the first two is  $H^0\mathcal{I}_{\{x_1, x_2\}}(2)$ . By Lemma 2 of 7.5.2,

$$\dim H^0\mathcal{I}_{\{x_1, x_2\}}(2) \cap H^0\mathcal{I}_3(2) = 2.$$

$\square$

#### 7.5.4. Proof of claim F:

a) We choose an isomorphism  $\bar{H}_3 \cong k^3$  and

$$\bar{\omega} = \begin{pmatrix} \eta_1 & 0 & 0 \\ 0 & \eta_2 & 0 \\ 0 & 0 & \eta_3 \end{pmatrix}$$

where  $\eta_i \in \Lambda^2 V^*$  are indecomposable. Let  $\mathcal{E}_i = \mathcal{E}_{\eta_i}$  be the associated nc-bundles such that

$$\mathcal{E} = \mathcal{E}_{\bar{\omega}} = \mathcal{E}_1 \oplus \mathcal{E}_2 \oplus \mathcal{E}_3.$$

By the definition of  $\tilde{X}_{\bar{\omega}}$  in 7.5 this fibre is isomorphic to  $\text{Hom}(k^3, \Lambda^2 V^*)$  and we have a surjective linear map

$$\tilde{X}_{\bar{\omega}} \rightarrow \text{Hom}(\mathcal{E}_1 \oplus \mathcal{E}_2 \oplus \mathcal{E}_3, \mathcal{O}(1))$$



by

$$(\alpha_1, \alpha_2, \alpha_3) \mapsto (\sigma_1(\alpha_1), \sigma_2(\alpha_2), \sigma_3(\alpha_3)).$$

The open set  $\tilde{X}_{\bar{\omega}}^0$  in  $\tilde{X}_{\bar{\omega}}$  is then the inverse image of the set of those  $(\sigma_1, \sigma_2, \sigma_3)$ ,  $\mathcal{E}_i \xrightarrow{\sigma_i} \mathcal{O}(1)$ , such that for the corresponding ideal sheaves  $\mathcal{I}_i$  we have

$$H^0\mathcal{I}_1(2) + H^0\mathcal{I}_2(2) + H^0\mathcal{I}_3(2) = H^0\mathcal{O}(2). \quad (*)$$

We put  $W_i = \text{Hom}(\mathcal{E}_i, \mathcal{O}(1))$  and let

$$Z \subset \mathbb{P}(W_1 \oplus W_2 \oplus W_3)$$

be the closed subvariety of points  $\langle \sigma_1, \sigma_2, \sigma_3 \rangle$  for which  $(*)$  is not satisfied. Then  $\text{codim } Z \geq 2$  implies claim F for the chosen  $\bar{\omega}$ .

b) In order to prove  $\text{codim } Z \geq 2$  we consider the natural projection

$$\mathbb{P}(W_1 \oplus W_2 \oplus W_3) \setminus Z' \xrightarrow{\pi} \mathbb{P}W_1 \times \mathbb{P}W_2 \times \mathbb{P}W_3$$

where  $Z'$  is the subvariety of points  $\langle \sigma_1, \sigma_2, \sigma_3 \rangle$  with at least one component equal to 0. It is easy to see that  $\pi$  is a principal bundle with fibre and group  $(k^*)^3/k^*$ . We have  $Z' \subset Z$  and  $Z \setminus Z'$  is invariant under  $(k^*)^3/k^*$ . Then  $Y = \pi(Z \setminus Z')$  is closed and  $Z \setminus Z' = \pi^{-1}(Y)$ . Now  $Y$  is the subvariety of triples  $(\langle \sigma_1 \rangle, \langle \sigma_2 \rangle, \langle \sigma_3 \rangle)$  for which  $(*)$  is not satisfied for the images  $\mathcal{I}_\nu(1) = \text{Im } \sigma_\nu$ . Claim F will be proved if  $\text{codim } Y \geq 2$ . This follows now from

c) **Proposition:** *Let  $\langle \eta_1 \rangle, \langle \eta_2 \rangle, \langle \eta_3 \rangle \in \mathbb{P}\Lambda^2 V^* \setminus G$  be in general position (not co-linear). Then the subvariety  $Y$  has codimension  $\geq 2$ .*

*Proof.* We let  $y_i$  denote the points  $\langle \eta_i \rangle$  in  $\mathbb{P}\Lambda^2 V$ .

(i) Let  $H(y_2)$  be the polar hyperplane of  $y_2$  with respect to the quadric  $G$ . There is point  $l \in H(y_2) \cap G$  such that the tangent hyperplane  $T_l G$  doesn't contain  $y_3$ . Denote by

$$C(l) = G \cap T_l(G)$$

the cone of lines in  $\mathbb{P}V$  meeting  $l$ . Then  $C(l) \cap H(y_2)$  has codimension 2.

(ii) Choose any  $l_{11} \in H(y_2) \cap G$  with  $y_3 \notin T_{l_{11}} G$  and such that the line  $g_1 = \overline{y_1, l_{11}}$  meets  $G$  in two different points  $l_{12} \neq l_{11}$ . We have  $l_{11}, y_2 \in T_{l_{11}} G$  because  $l_{11} \in H(y_2)$ . Next we choose a line  $g_2$  in  $T_{l_{11}} G$  through  $y_2$  which meets  $G$  in two different points  $l_{21}, l_{22}$ . These lines belong to the cone  $C(l_{11})$  and thus each meets  $l_{11}$  in a point  $x_1$  and  $x_2$ . We may assume that  $g_2 \cap C(l_{12}) = \emptyset$ , such that  $l_{21}$  and  $l_{22}$  don't meet  $l_{12}$ . By this choice the conditions of 7.5.3, Lemma 3, are satisfied, and thus

$$H^0\mathcal{I}_{g_1}(2) + H^0\mathcal{I}_{g_2}(2) = H^0\mathcal{I}_{\{x_1, x_2\}}(2)$$

is 8-dimensional, where  $\mathcal{I}_{g_\nu}$  is the ideal corresponding to the line  $g_\nu$  through  $y_\nu$ .

(iii) Let now  $P(x_\nu) \subset G$  be the  $\alpha$ -plane of all lines through  $x_\nu$  and let

$$S = P(x_1) \cup P(x_2) \cup (C(l_{11}) \cap H(y_3)).$$

Because  $y_3 \notin T_{l_{11}}G$ ,  $C(l_{11}) \cap H(y_3)$  is 2-dimensional and hence  $S$  is 2-dimensional. Let

$$\mathbb{P}\Lambda^2V \setminus \{y_3\} \xrightarrow{\pi} \mathbb{P}(\Lambda^2V/y_3)$$

be the central projection. We have  $y_3 \notin S$ . Then  $\pi(S)$  has dimension 2 and there is a line  $L \subset \mathbb{P}(\Lambda^2V/y_3)$  which doesn't meet  $\pi(S)$ . Let  $a_3 \in L$  be any point and let  $g_3$  denote the line through  $y_3$  in  $\mathbb{P}\Lambda^2V$  given by  $\pi^{-1}(a_3)$ . If  $a_3 \notin \pi(H(y_3) \cap G)$ , the branch locus of  $\pi|_G$ , then  $g_3 \cap G$  consists of two different points  $l_{31}, l_{32}$  which don't meet  $x_1, x_2$ . Suppose that  $l_{31}$  meets  $l_{11}$ , i.e.  $l_{31} \in C(l_{11}) \subset T_{l_{11}}G$ . Then  $l_{32} \notin T_{l_{11}}G$  because otherwise also  $y_3 \in T_{l_{11}}G$ , which had been excluded by the choice of  $l_{11}$ . By the corollary in 7.5.3  $H^0\mathcal{I}_{g_3}(2)$  intersects  $H^0\mathcal{I}_{\{x_1, x_2\}}(2)$  in dimension 2. Then for  $\langle \alpha_\nu \rangle \in g_\nu, \nu = 1, 2, 3$ , we have a point  $(\langle \alpha_1 \rangle, \langle \alpha_2 \rangle, \langle \alpha_3 \rangle) \in \mathbb{P}W_1 \times \mathbb{P}W_2 \times \mathbb{P}W_3 \setminus Y$ . If however,  $a_3 \in \pi(H(y_3) \cap G)$ , then  $g_3$  is tangent to  $G$  at some  $l_0$  and  $\mathcal{I}_{g_3}$  is a twisted double structure on  $l_0$ . In this case  $l_0 \notin C(l_{11})$  because  $l_0 \in H(y_3)$ , and so  $l_0 \cap l_{11} = \emptyset$ . Again by the corollary  $H^0\mathcal{I}_{g_3}(2)$  and  $H^0\mathcal{I}_{\{x_1, x_2\}}(2)$  intersect in dimension 2.

(iv) Let now  $a'_3, a''_3 \in L$  be any two points,  $g'_3, g''_3$  the lines in  $\mathbb{P}\Lambda^2V$  through  $y_3$ , defined by  $\pi^{-1}(a'_3), \pi^{-1}(a''_3)$  and let  $\langle \alpha_1 \rangle \in g_1 \setminus \{y_1\}, \langle \alpha_2 \rangle \in g_2 \setminus \{y_2\}, \langle \alpha'_3 \rangle \in g'_3 \setminus \{y_3\}, \langle \alpha''_3 \rangle \in g''_3 \setminus \{y_3\}$ . Then the points

$$(\langle \alpha_1 \rangle, \langle \alpha_2 \rangle, \langle s\alpha'_3 + t\alpha''_3 \rangle) \in \mathbb{P}W_1 \times \mathbb{P}W_2 \times \mathbb{P}W_3$$

do not belong to  $Y$  for any  $(s, t) \neq (0, 0)$ . We thus have a complete curve in the product space not meeting  $Y$ , i.e.  $\text{codim} Y \geq 2$ . This completes the proof of claim F and with that the proof of Lemma 7.2 and finally the proof of theorem 7.3.  $\square$

## 8. SMOOTHNESS OF $MI(5)$

The induction step  $\mathcal{E}_\omega \rightsquigarrow \mathcal{E}_{\bar{\omega}}$  will be used in order to prove that  $H^2S^2\mathcal{E}_\omega = 0$  follows from  $H^2S^2\mathcal{E}_{\bar{\omega}} = 0$  even if  $h^1\mathcal{E}_{\bar{\omega}}(1) = 1$ .

As before we use the following notation.  $\mathcal{E}$  will denote a 5-instanton and  $\bar{\mathcal{E}}$  the  $(4, 4)$ -instanton obtained from a general  $\xi \in H_5^* \cong H^1\mathcal{E}(-1)$  by the process  $\omega \mapsto \bar{\omega} = \text{res}_\xi \omega$ , such that  $\mathcal{E} = \mathcal{E}_\omega$  and  $\bar{\mathcal{E}} = \mathcal{E}_{\bar{\omega}}$ , see 4.5.

By Proposition 4.5 we may assume that there is a second element  $\eta \in H^1\mathcal{E}(-1)$ , linear independent of  $\xi$ , such that the multiplication map  $\langle \xi, \eta \rangle \otimes V^* \rightarrow H^1\mathcal{E}$  is injective (in fact an isomorphism). Let  $\bar{H} = \text{Ker}(\xi) \subset H_5$  and let  $\bar{\eta} \in \bar{H}_4^*$  be the image of  $\eta$ . If  $\bar{\omega} = \text{res}_\xi \omega$ , then  $\bar{\omega} \in M(4, 4)_{\bar{\eta}}$ , and we have  $H^2(S^2\bar{\mathcal{E}}) = 0$ , see 3.2 in case  $n = 4$ . Now  $\mathcal{E}$  appears as the cohomology of the monad

$$0 \rightarrow \mathcal{O}_{\mathbb{P}}(-1) \rightarrow \bar{\mathcal{E}} \rightarrow \mathcal{O}_{\mathbb{P}}(1) \rightarrow 0$$

resulting from  $\xi$ . Then  $S^2\mathcal{E}$  appears as the cohomology of the derived monad

$$0 \rightarrow \bar{\mathcal{E}}(-1) \rightarrow S^2\bar{\mathcal{E}} \oplus \mathcal{O} \rightarrow \bar{\mathcal{E}}(1) \rightarrow 0,$$

see proof of 2.8.1 and 3.2. Because of the vanishing conditions for instantons, we obtain an exact sequence

$$H^1(S^2\bar{\mathcal{E}}) \rightarrow H^1\bar{\mathcal{E}}(1) \rightarrow H^2(S^2\mathcal{E}) \rightarrow H^2(S^2\bar{\mathcal{E}}) = 0.$$

By Proposition 6.1 we may assume that  $h^1\bar{\mathcal{E}}(1) \leq 1$ . In what follows, we shall, practically, describe the kernel of the map  $H^2(\bar{\mathcal{E}}(1))^* \rightarrow H^1(S^2\bar{\mathcal{E}})^*$  and show that it is zero. Then it follows that  $H^2S^2\mathcal{E} = 0$ . In order to do that, we first describe the dual of  $H^1\bar{\mathcal{E}}(1)$  in terms of its defining tensor  $\bar{\omega}$ , which can be done for arbitrary  $n$ .

**8.1. Lemma:** *Let  $H$  be an  $n$ -dimensional vector space, let  $\omega \in S^2H^* \otimes \Lambda^2V^*$  be non-degenerate and let  $\mathcal{E} = \mathcal{E}_\omega$ , and interpret the elements of  $H \otimes S^2V$  as linear maps  $H^* \otimes V^* \rightarrow V$  which are selfdual with respect to  $V$ . Then*

- (a)  $H^1(\mathcal{E}(1))^* \cong \{\gamma \in H \otimes S^2V \mid \gamma \circ \omega = 0\}$
- (b) *If  $P = \mathbb{P}W$  is a plane in  $\mathbb{P}_3$ ,  $W \subset V$ , then  $H^1(\mathcal{E}_P(1))^* \cong \{\gamma \in H \otimes S^2V \mid \gamma \circ \omega = 0 \text{ and } \text{Im } \gamma \subset W\}$*

*Proof.* a) From the selfdual monad associated to  $\omega$ , one derives the exact sequence

$$N \otimes V^* \rightarrow H^* \otimes S^2V^* \rightarrow H^1\mathcal{E}(1) \rightarrow 0$$

and its dual

$$0 \rightarrow H^1(\mathcal{E}(1))^* \rightarrow H \otimes S^2V \rightarrow N^* \otimes V.$$

Interpreting the elements of  $N^* \otimes V$  as linear maps  $N \rightarrow V$  and the elements of  $H \otimes S^2V$  as linear maps  $H^* \otimes V^* \xrightarrow{\gamma} V \cong H_1 \otimes V$ , the homomorphism in the last sequence can be described by  $\gamma \mapsto \gamma \circ u$ , where  $u : N \hookrightarrow H^* \otimes V^*$ .

Since  $N = \text{Im}(\omega)$ , it follows that  $\gamma \circ u = 0$  if and only if  $\gamma \circ \omega = 0$ .

b) If  $\gamma \in H \otimes S^2V$  and  $\text{Im}(\gamma) \subset W$ , then  $\gamma \in H \otimes S^2W$ , i.e.  $\gamma$  factorizes as

$$\begin{array}{ccc} H^* \otimes V^* & \xrightarrow{\gamma} & V \\ \downarrow & & \uparrow \\ H^* \otimes W^* & \xrightarrow{\gamma_0} & W \end{array}$$

for some  $\gamma_0$  which is selfdual with respect to  $W$ . Now restrict the monad of  $\omega$  to  $P$  and use the same argument as in a).  $\square$

**8.2. Proposition:** *Any  $(5, 2)$ -instanton  $\mathcal{E}$  on  $\mathbb{P}_3$  satisfies  $H^2S^2\mathcal{E} = 0$ .*

*Proof.* a) Let  $\omega \in M(5, 2)$  define  $\mathcal{E} = \mathcal{E}_\omega$ . As before, we denote by  $\bar{\mathcal{E}}$  the bundle  $\mathcal{E}_{\bar{\omega}}$  defined by  $\bar{\omega} = \text{res}_\xi(\omega)$  for a general  $\xi \in H^*$ . We may assume that  $h^1\bar{\mathcal{E}}(1) \leq 1$  by Proposition 6.1. As mentioned above, we may also assume that  $H^2S^2\bar{\mathcal{E}} = 0$ .

b) Let  $H_1 \oplus \bar{H} = H$  be a decomposition defined by  $\xi$  with  $\bar{H} = \text{Ker}(\xi)$ . Then

$$(H_1 \oplus \bar{H}) \otimes V \xrightarrow{\omega} (H_1^* \oplus \bar{H}^*) \otimes V^*$$

decomposes as

$$\omega = \left( \begin{array}{c|c} e \circ \bar{\varphi} \circ e^* & e \circ \bar{\varphi} \circ \bar{u}^* \\ \hline \bar{u} \circ \bar{\varphi} \circ e^* & \bar{\omega} \end{array} \right)$$

where  $\bar{N} \xrightarrow{e} H_1^* \otimes V^*$  is a linear map and where

$$\begin{array}{ccc} \bar{H} \otimes V & \xrightarrow{\bar{\omega}} & \bar{H}^* \otimes V^* \\ \downarrow \bar{u}^* & & \uparrow \bar{u} \\ \bar{N} & \xrightarrow[\cong]{\bar{\varphi}} & \bar{N} \end{array}$$

is the decomposition of  $\bar{\omega} = \bar{u} \circ \bar{\varphi} \circ \bar{u}^*$ . Note that  $\omega$  is symmetric, resp. skew with respect to  $H$ , resp.  $V$ . Similarly, any  $\sigma \in \Lambda^2 H \otimes S^2 V$  can be written as a linear operator

$$(H_1^* \oplus \bar{H}^*) \otimes V^* \xrightarrow{\sigma} (H_1 \oplus \bar{H}) \otimes V$$

and decomposed as

$$\sigma = \left( \begin{array}{c|c} 0 & \gamma \\ \hline -\gamma^* & \bar{\sigma} \end{array} \right)$$

with  $\gamma \in \bar{H} \otimes S^2 V$  because  $\sigma$  is skew with respect to  $H$ .

c) We are now using the exact sequence

$$0 \rightarrow H^2(S^2\mathcal{E})^* \rightarrow \Lambda^2 H \otimes S^2 V \rightarrow N^* \otimes H \otimes V$$

of Remark 2.10, the homomorphism being  $\sigma \mapsto \sigma \circ \omega$ . By the above decomposition the condition  $\sigma \circ \omega = 0$  is equivalent to the two conditions

$$\gamma \circ \bar{u} = 0 \text{ and } \gamma^* \circ e = \bar{\sigma} \circ \bar{u}.$$

In order to show that  $\sigma \circ \omega = 0$  implies  $\sigma = 0$ , it is now sufficient to prove that the two conditions on  $\gamma$  imply that  $\gamma = 0$ . For then, also,  $\bar{\sigma} \circ \bar{u} = 0$  or  $\bar{\sigma} \circ \bar{\omega} = 0$ , and then  $\bar{\sigma} = 0$ , because  $H^2S^2\bar{\mathcal{E}} = 0$ , using the same sequence for  $\bar{\mathcal{E}}$ .

d) The first condition  $\gamma \circ \bar{u} = 0$  or  $\gamma \circ \bar{\omega} = 0$  means that

$$\gamma \in H^1(\bar{\mathcal{E}}(1))^* \subset \bar{H} \otimes S^2 V$$

by Lemma 8.1, (a). If  $H^1\bar{\mathcal{E}}(1) = 0$ , there is nothing to prove. So we may assume that  $\gamma \neq 0$ . By Lemma 6.3 we may assume that  $H^1\bar{\mathcal{E}}_P(1) = 0$  for any plane  $P = \mathbb{P}W$  in

$\mathbb{P}V$ . Then, by Lemma 8.1, (b),  $\text{Im}(\gamma) \not\subset W$  for any 3-dimensional subspace  $W \subset V$ . Therefore,

$$\bar{H}^* \otimes V^* \xrightarrow{\gamma} H_1 \otimes V$$

is surjective. Because  $\gamma \circ \bar{u} = 0$  and  $\dim \bar{N} = 12$ , we obtain the exact diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bar{N} & \longrightarrow & \bar{H}^* \otimes V^* & \xrightarrow{\gamma} & H_1 \otimes V \longrightarrow 0 \\ & & \uparrow \cong & & \uparrow & & \uparrow \\ 0 & \longrightarrow & N & \longrightarrow & H^* \otimes V^* & \longrightarrow & Q \longrightarrow 0 \\ & & & & \uparrow & & \\ & & & & \xi \otimes V^* & & \\ & & & & \uparrow & & \\ & & & & 0. & & \end{array}$$

Note that the induced projection induces an isomorphism between  $N$  and  $\bar{N}$ , because  $\omega$  and  $\bar{\omega}$  have the same rank 12. It follows from the decomposition of  $\omega$  that any vector of  $N$  can be written uniquely as

$$e(\theta) + \theta$$

with  $\theta \in \bar{N} \subset \bar{H}^* \otimes V^*$ , with the isomorphism given by  $e(\theta) + \theta \leftrightarrow \theta$ , and  $\text{Im}(e) \subset \xi \otimes V^*$ .

e) We are now going to show that the condition  $\gamma^* \circ e = \bar{\sigma} \circ \bar{u}$  implies a contradiction. For that we choose a 2-dimensional subspace  $K \subset H^*$  containing  $\xi$ , using Lemma 4.7, such that  $N \cap (K \otimes V^*)$  contains no non-zero decomposable vector. Let  $\xi, \xi'$  be the basis of  $K$ ,  $\xi' \in \bar{H}^*$ . Then there are two independent linear forms  $z, z' \in V^*$  such that  $N \cap (K \otimes V^*)$  is spanned by

$$\xi \otimes z + \xi' \otimes z'.$$

Because  $\xi \otimes z \mapsto 0$  under  $N \rightarrow \bar{N}$ , we have with  $\theta = \xi' \otimes z'$  that

$$\xi \otimes z + \xi' \otimes z' = e(\xi' \otimes z') + \xi' \otimes z'$$

or  $e(\xi' \otimes z') = \xi \otimes z$ . Therefore,  $\gamma^*(\xi \otimes z) = \bar{\sigma}(\xi' \otimes z')$ .

Because  $\bar{\sigma} \in \Lambda^2 \bar{H} \otimes S^2 V$  is skew with respect to  $\bar{H}$ , it follows  $\bar{\sigma}(\eta \otimes w) \in \text{Ker}(\eta) \otimes V$  under  $\bar{H}^* \otimes V^* \xrightarrow{\bar{\sigma}} \bar{H} \otimes V$  for any  $\eta \in \bar{H}^*$  and any  $w \in V^*$ . It follows that  $(\xi' \otimes \text{id}) \circ \gamma^*(\xi' \otimes z) = 0$ , and dually, that  $\gamma(\xi' \otimes z) = 0$ , by considering the diagrams

$$\begin{array}{ccc} H_1^* \otimes V^* & \xrightarrow{\gamma^*} & \bar{H} \otimes V \\ \uparrow & & \downarrow \xi' \otimes \text{id} \\ \langle \xi \otimes z \rangle & \xrightarrow{0} & V \end{array} \quad \begin{array}{ccc} H_1 \otimes V & \xleftarrow{\gamma^*} & \bar{H}^* \otimes V^* \\ \downarrow \xi \otimes z & & \uparrow \\ k & \xleftarrow{0} & \xi' \otimes V^* \end{array}$$

and using that  $\gamma$  is symmetric with respect to  $V$ .

Now  $\xi' \otimes z \in \bar{N}$  and, because  $e$  has its image in  $\xi \otimes V^*$ , there is a form  $z'' \in V^*$  with  $e(\xi' \otimes z) = \xi \otimes z''$ . Then

$$\xi \otimes z'' + \xi' \otimes z \in N \cap (K \otimes V^*)$$

is a second vector which is independent of  $\xi \otimes z + \xi' \otimes z'$ , a contradiction to the choice of  $K$ . Therefore, the condition  $\gamma^* \circ e = \bar{\sigma} \circ \bar{u}$  for  $\gamma \in H^1(\bar{\mathcal{E}}(1))^*$  implies  $\gamma = 0$ , which proves Proposition 8.2.  $\square$

## REFERENCES

- [1] **M. Atiyah - N. Hitchin - V.G. Drinfeld - Y. Manin**, Construction of Instantons, Phys. Letters 65 A, 185-187, 1978
- [2] **W. Barth**, Some properties of stable rank-2 vector bundles on  $\mathbb{P}_n$ , Math. Ann. 226, 125-150, 1977
- [3] **W. Barth**, Irreducibility of the Space of Mathematical Instanton Bundles with Rank 2 and  $c_2 = 4$ , Math. Ann. 258, 81-106, 1981
- [4] **W. Barth - K. Hulek**, Monads and moduli of vector bundles, manuscripta math. 25, 323-347, 1978
- [5] **A. Beilinson**, Coherent sheaves on  $\mathbb{P}^N$  and problems of linear algebra, Funct. Anal. Appl. 12, 214-216, 1978
- [6] **W. Böhmer - G. Trautmann**, Special Instanton Bundles and Poncelet Curves, in Singularities, Representation of Algebras, and Vector Bundles, Lecture Notes in Mathematics 1273, Springer-Verlag 1987
- [7] **I. Coandă**, On Barth's restriction theorem, Journ. reine u. angew. Mathematik 428, 97-110, 1992
- [8] **I. Coandă**, Restriction theorems for vector bundles via the method of Roth and Strano, Journ. reine u. angew. Mathematik 487, 1-25, 1997
- [9] **L. Costa - G. Ottaviani**, Nondegenerate multidimensional matrices and instanton bundles, preprint math.AG/0103078
- [10] **G. Ellingsrud - S.A. Strømme**, Stable rank-2 vector bundles on  $\mathbb{P}_3$  with  $c_1 = 0$  and  $c_2 = 3$ , Math. Ann. 255, 123-135, 1981
- [11] **A. Grothendieck - J. Dieudonné**, Eléments de Géométrie Algébrique I, Springer-Verlag 1971
- [12] **R. Hartshorne**, Stable vector bundles of rank 2 on  $\mathbb{P}_3$ , Math. Ann. 238, 229-280, 1978
- [13] **A. Hirschowitz - M.S. Narasimhan**, Fibré de 'tHooft spéciaux et applications, Proc. Nice Conf. 1981, Birkhäuser 1982
- [14] **S.L. Kleiman**, Relative duality for quasi-coherent sheaves, Compositio Mathematica 41, 39-60, 1980
- [15] **P.I. Katsylo - G. Ottaviani**, Regularity of the Moduli Space of Instanton Bundles  $MI_{\mathbb{P}_3}(5)$ , math.AG/9911184
- [16] **D. Luna**, Slices étales, Bull. Soc. Math. France 33, 81-105, 1973
- [17] **M. Maruyama**, Moduli of stable sheaves I, J. Math. Kyoto Univ. 17, 91-126, 1977
- [18] **M. Maruyama**, Moduli of stable sheaves II, J. Math. Kyoto Univ. 18, 557-614, 1978
- [19] **Th. Nüßler - G. Trautmann**, Multiple Koszul structures on lines and instanton bundles, Intern. Journ. of Math. 3, 373-388, 1994
- [20] **C. Okonek, M. Schneider, H. Spindler**, Vector Bundles on Complex Projective Spaces, Birkhäuser 1980

- [21] **J. LePotier**, Sur l'espace de modules des fibrés de Yang et Mills, in Mathématique et Physique, Séminaire de l'Ecole Normale Supérieure 1979-1982, Birkhäuser 1983
- [22] **P. Rao**, Mathematical instantons with maximal order jumping lines, Pacific Journ. of Mathem. 178, 331-344, 1997
- [23] **I.R. Shafarevich**, Basic algebraic geometry, 2nd ed., Springer Verlag 1994
- [24] **M. Skiti**, Sur une famille de fibrés instantons, Math. Z. 225, 373-294, 1997
- [25] **A.N. Tyurin**, On the Superposition of Mathematical Instantons II, in Arithmetic and Geometry, Progress in Mathematics 36, Birkhäuser 1983
- [26] **A.N. Tyurin**, The structure of the variety of pairs of commuting pencils of symmetric matrices, Izv. Akad. Nauk SSSR, Ser. Matem. Tom 46, no. 2 (1982), English translation: Math. USSR Izvestiya, Vol. 20, no. 2 (1983), 391-410
- [27] **J.L. Verdier**, Instantons, in: Les équations de Yang-Mills, Séminaire E.N.S. 1977/78, Exposé VII, Astérisque 71-72, 1980

INSTITUTE OF MATHEMATICS, OF THE ROMANIAN ACADEMY, P.O. Box 1-764  
RO-70700 BUCHAREST, ROMANIA

*E-mail address:* `Iustin.Coanda@imar.ro`

DEPARTMENT OF MATHEMATICS, STATE PEDAGOGICAL UNIVERSITY, RESPUBLIKANSKAYA STR. 108  
150 000 YAROSLAVL, RUSSIA

*E-mail address:* `alexandr@tikhomir.yaroslavl.su`

UNIVERSITÄT KAISERSLAUTERN, FACHBEREICH MATHEMATIK, ERWIN-SCHRÖDINGER-STRASSE  
D-67663 KAISERSLAUTERN

*E-mail address:* `trm@mathematik.uni-kl.de`